



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁵ : C12N 15/00, C07K 13/00 G01N 33/531, 33/68</p>	A1	<p>(11) International Publication Number: WO 92/01047</p> <p>(43) International Publication Date: 23 January 1992 (23.01.92)</p>															
<p>(21) International Application Number: PCT/GB91/01134</p> <p>(22) International Filing Date: 10 July 1991 (10.07.91)</p> <p>(30) Priority data:</p> <table style="width: 100%; border: none;"> <tr> <td style="width: 30%;">9015198.6</td> <td style="width: 30%;">10 July 1990 (10.07.90)</td> <td style="width: 40%;">GB</td> </tr> <tr> <td>9022845.3</td> <td>19 October 1990 (19.10.90)</td> <td>GB</td> </tr> <tr> <td>9024503.6</td> <td>12 November 1990 (12.11.90)</td> <td>GB</td> </tr> <tr> <td>9104744.9</td> <td>6 March 1991 (06.03.91)</td> <td>GB</td> </tr> <tr> <td>9110549.4</td> <td>15 May 1991 (15.05.91)</td> <td>GB</td> </tr> </table> <p>(71) Applicants (for all designated States except US): CAMBRIDGE ANTIBODY TECHNOLOGY LIMITED [GB/GB]; The Daly Research Laboratories, Babraham Hall, Cambridge, Cambridgeshire CB2 4AT (GB). MEDICAL RESEARCH COUNCIL [GB/GB]; 20 Park Crescent, London W1N 4AL (GB).</p> <p>(72) Inventors; and</p> <p>(75) Inventors/Applicants (for US only) : McCAFFERTY, John [GB/GB]; 32 Wakelin Avenue, Sawston, Cambridgeshire CB2 4DS (GB). POPE, Anthony, Richard [GB/GB]; 31 Kingston Street, Cambridge CB1 2NU (GB). JOHNSON, Kevin, Stuart [GB/GB]; 32 Holmehill, Godmanchester, Cambridgeshire PE18 8EX (GB). HOOGENDOORN, Hendricus, Renerus, Jacobus, Mattheus [NL/GB]; 25 Queensway, Cambridge CB2 2AY (GB). GRIFFITHS, Andrew, David [GB/GB]; 28 Lilac Court, Cherry Hinton Road, Cambridge CB1 4AY (GB). JACKSON, Ronald, Henry [GB/GB]; 31 Kingston Street, Cambridge CB1 2NU (GB). HOLLIGER, Kaspar,</p>			9015198.6	10 July 1990 (10.07.90)	GB	9022845.3	19 October 1990 (19.10.90)	GB	9024503.6	12 November 1990 (12.11.90)	GB	9104744.9	6 March 1991 (06.03.91)	GB	9110549.4	15 May 1991 (15.05.91)	GB
9015198.6	10 July 1990 (10.07.90)	GB															
9022845.3	19 October 1990 (19.10.90)	GB															
9024503.6	12 November 1990 (12.11.90)	GB															
9104744.9	6 March 1991 (06.03.91)	GB															
9110549.4	15 May 1991 (15.05.91)	GB															
<p>Philipp [CH/GB]; 80 Hobart Road, Cambridge CB1 3PT (GB). MARKS, James, David [US/GB]; 82 Mill End Road, Cherry Hinton, Cambridge CB1 4JP (GB). CLACKSON, Timothy, Piers [GB/GB]; 225 Mill Road, Cambridge CB1 3BE (GB). CHISWELL, David, John [GB/GB]; 1 Sandhill House, Middle Claydon, Buckingham MK18 2LD (GB). WINTER, Gregory, Paul [GB/GB]; 64 Cavendish Avenue, Cambridge CB1 4UT (GB). BONNERT, Timothy, Peter [GB/GB]; 77 Glisson Road, Cambridge CB1 2HG (GB).</p> <p>(74) Agent: MEWBURN, Ellis; 2 Cursitor Street, London EC4A 1BQ (GB).</p> <p>(81) Designated States: AT, AT (European patent), AU, BB, BE (European patent), BF (OAPI patent), BG, BJ (OAPI patent), BR, CA, CF (OAPI patent), CG (OAPI patent), CH, CH (European patent), CI (OAPI patent), CM (OAPI patent), CS, DE, DE (European patent), DK, DK (European patent), ES, ES (European patent), FI, FR (European patent), GA (OAPI patent), GB, GB (European patent), GN (OAPI patent), GR (European patent), HU, IT (European patent), JP, KP, KR, LK, LU, LU (European patent), MC, MG, ML (OAPI patent), MN, MR (OAPI patent), MW, NL, NL (European patent), NO, PL, RO, SD, SE, SE (European patent), SN (OAPI patent), SU, TD (OAPI patent), TG (OAPI patent), US.</p> <p>Published With international search report.</p>																	
<p>(54) Title: METHODS FOR PRODUCING MEMBERS OF SPECIFIC BINDING PAIRS</p> <p>(57) Abstract</p> <p>A member of a specific binding pair (sbp) is identified by expressing DNA encoding a genetically diverse population of such sbp members in recombinant host cells in which the sbp members are displayed in functional form at the surface of a secreted recombinant genetic display package (rgdp) containing DNA encoding the sbp member or a polypeptide component thereof, by virtue of the sbp member or a polypeptide component thereof being expressed as a fusion with a capsid component of the rgdp. The displayed sbps may be selected by affinity with a complementary sbp member, and the DNA recovered from selected rgdps for expression of the selected sbp members. Antibody sbp members may be thus obtained, with the different chains thereof expressed, one fused to the capsid component and the other in free form for association with the fusion partner polypeptide. A phagemid may be used as an expression vector, with said capsid fusion helping to package the phagemid DNA. Using this method libraries of DNA encoding respective chains of such multimeric sbp members may be combined, thereby obtaining a much greater genetic diversity in the sbp members than could easily be obtained by conventional methods.</p>																	
<div style="display: flex; justify-content: space-around; align-items: flex-start;"> <div style="text-align: center;"> <p>Fab</p> </div> <div style="text-align: center;"> </div> <div style="text-align: center;"> <p>scFv</p> </div> </div>																	

+ DESIGNATIONS OF "SU"

It is under examination in which parts of the former Soviet Union the designation of the Soviet Union has effect.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland			SN	Senegal
CI	Côte d'Ivoire	KR	Republic of Korea	SU	Soviet Union
CM	Cameroon	LI	Liechtenstein	TD	Chad
CS	Czechoslovakia	LK	Sri Lanka	TC	Togo
DE	Germany	LU	Luxembourg	US	United States of America
DK	Denmark	MC	Monaco		

METHODS FOR PRODUCING MEMBERS OF SPECIFIC BINDING PAIRS

5 The present invention relates to methods for producing members of specific binding pairs. The present invention also relates to the biological binding molecules produced by these methods.

10 Owing to their high specificity for a given antigen, the advent of monoclonal antibodies (Kohler, G. and Milstein C; 1975 Nature 256: 495) represented a significant technical break-through with important consequences both scientifically and commercially.

15 Monoclonal antibodies are traditionally made by establishing an immortal mammalian cell line which is derived from a single immunoglobulin producing cell secreting one form of a biologically functional antibody molecule with a particular specificity. Because the antibody-secreting mammalian cell line is immortal, the characteristics of the antibody are reproducible from batch to batch. The key properties of monoclonal antibodies are their specificity for a particular antigen and the reproducibility with which they can be manufactured.

20 Structurally, the simplest antibody (IgG) comprises four polypeptide chains, two heavy (H) chains and two light (L) chains inter-connected by disulphide bonds (see figure 1). The light chains exist in two distinct forms called kappa (K) and lambda (λ). Each chain has a constant region (C) and a variable region (V). Each chain is organized into a series of domains. The light chains have two domains, corresponding to the C region and the other to the V region. The heavy chains have four domains, one corresponding to the V region and three domains (1,2 and 3) in the C region. The antibody has two arms (each arm being a Fab region), each of which has a VL and a VH region associated with each other. 25 It is this pair of V regions (VL and VH) that differ from one antibody to another (owing to amino acid sequence variations), and which together are responsible for recognising the antigen and providing an antigen binding site (ABS). In even more detail, each V region is made up from three complementarity determining regions (CDR) separated by four framework regions (FR). The CDR's are the most variable part of the variable regions, and they perform the critical antigen binding function. The CDR regions are derived from many potential germ line sequences via a complex process involving recombination, mutation and selection. 30 35 40 45

46 It has been shown that the function of binding antigens can be performed by fragments of a whole antibody. Example binding fragments are (i) the Fab fragment consisting of the VL, VH, CL and CH1 domains; (ii) the Fd fragment consisting of the VH and CH1 domains; (iii) the Fv fragment consisting of the VL and VH domains of a single arm of an antibody, (iv) the dAb fragment (Ward, E.S. et al., Nature 341, 544-546 (1989) which consists of a VH domain; (v) isolated CDR 50

regions; and (vi) $F(ab')_2$ fragments, a bivalent fragment comprising two Fab fragments linked by a disulphide bridge at the hinge region.

Although the two domains of the Fv fragment are coded for by separate genes, it has proved possible to make a synthetic linker that enables them to be made as a single protein chain (known as single chain Fv (scFv); Bird, R.E. et al., *Science* 242, 423-426 (1988) Huston, J.S. et al., *Proc. Natl. Acad. Sci., USA* 85, 5879-5883 (1988)) by recombinant methods. These scFv fragments were assembled from genes from monoclonals that had been previously isolated. In this application, the applicants describe a process to assemble scFv fragments from VH and VL domains that are not part of an antibody that has been previously isolated.

Whilst monoclonal antibodies, their fragments and derivatives have been enormously advantageous, there are nevertheless a number of limitations associated with them.

Firstly, the therapeutic applications of monoclonal antibodies produced by human immortal cell lines holds great promise for the treatment of a wide range of diseases (Clinical Applications of Monoclonal Antibodies. Edited by E. S. Lennox. British Medical Bulletin 1984. Publishers Churchill Livingstone). Unfortunately, immortal antibody-producing human cell lines are very difficult to establish and they give low yields of antibody (approximately 1 $\mu\text{g/ml}$). In contrast, equivalent rodent cell lines yield high amounts of antibody (approximately 100 $\mu\text{g/ml}$). However, the repeated administration of these foreign rodent proteins to humans can lead to harmful hypersensitivity reactions. In the main therefore, these rodent-derived monoclonal antibodies have limited therapeutic use.

Secondly, a key aspect in the isolation of monoclonal antibodies is how many different clones of antibody producing cells with different specificities, can be practically established and sampled compared to how many theoretically need to be sampled in order to isolate a cell producing antibody with the desired specificity characteristics (Milstein, C., Royal Soc. Croonian Lecture, *Proc. R. Soc. London B.* 239; 1-16, (1990)). For example, the number of different specificities expressed at any one time by lymphocytes of the murine immune system is thought to be approximately 10^7 and this is only a small proportion of the potential repertoire of specificities. However, during the isolation of a typical antibody producing cell with a desired specificity, the investigator is only able to sample 10^3 to 10^4 individual specificities. The problem is worse in the human, where one has approximately 10^{12} lymphocyte specificities, with the limitation on sampling of 10^3 or 10^4 remaining.

This problem has been alleviated to some extent in laboratory animals by the use of immunisation regimes. Thus, where one wants to produce monoclonal antibodies having a specificity against a particular epitope, an animal

is immunised with an immunogen expressing that epitope. The animal will then mount an immune response against the immunogen and there will be a proliferation of lymphocytes which have specificity against the epitope. Owing to this proliferation of lymphocytes with the desired specificity, it becomes easier to detect them in the sampling procedure. However, this approach is not successful in all cases, as a suitable immunogen may not be available. Furthermore, where one wants to produce human monoclonal antibodies (eg for therapeutic administration as previously discussed), such an approach is not practically, or ethically, feasible.

In the last few years, these problems have in part, been addressed by the application of recombinant DNA methods to the isolation and production of e.g. antibodies and fragments of antibodies with antigen binding ability, in bacteria such as *E.coli*.

This simple substitution of immortalised cells with bacterial cells as the 'factory', considerably simplifies procedures for preparing large amounts of binding molecules. Furthermore, a recombinant production system allows scope for producing tailor-made antibodies and fragments thereof. For example, it is possible to produce chimaeric molecules with new combinations of binding and effector functions, humanised antibodies (e.g. murine variable regions combined with human constant domains or murine-antibody CDRs grafted onto a human FR) and novel antigen-binding molecules. Furthermore, the use of polymerase chain reaction (PCR) amplification (Saiki, R.K., et al., *Science* **239**, 487-491 (1988)) to isolate antibody producing sequences from cells (e.g. hybridomas and B cells) has great potential for speeding up the timescale under which specificities can be isolated. Amplified VH and VL genes are cloned directly into vectors for expression in bacteria or mammalian cells (Orlandi, R., et al., 1989, *Proc. Natl. Acad. Sci., USA* **86**, 3833-3837; Ward, E.S., et al., 1989 *supra*; Larrick, J.W., et al., 1989, *Biochem. Biophys. Res. Commun.* **160**, 1250-1255; Sastry, L. et al., 1989, *Proc. Natl. Acad. Sci., USA*, **86**, 5728-5732). Soluble antibody fragments secreted from bacteria are then screened for binding activities.

However, like the production system based upon immortalised cells, the recombinant production system still suffers from the selection problems previously discussed and therefore relies on animal immunization to increase the proportion of cells with desired specificity. Furthermore, some of these techniques can exacerbate the screening problems. For example, large separate H and L chain libraries have been produced from immunized mice and combined together in a random combinatorial manner prior to screening (Huse, W.D. et al., 1989, *Science* **246**, 1275-1281, W090/14443; W090/14424 and W090/14430). Crucially however, the information held within each cell, namely the original pairing of one L chain with one H chain, is lost. This loses some, of the advantage gained by using immunization protocols in the animal. Currently, only libraries derived from single VH domains (dAbs; Ward, E.S., et al., 1989,

supra.) do not suffer this drawback. However, because not all antibody VH domains are capable of binding antigen, more have to be screened. In addition, the problem of directly screening many different specificities in prokaryotes remains to be solved.

Thus, there is a need for a screening system which ameliorates or overcomes one or more of the above or other problems. The ideal system would allow the sampling of very large numbers of specificities (eg 10^6 and higher), rapid sorting at each cloning round, and rapid transfer of the genetic material coding for the binding molecule from one stage of the production process, to the next stage.

The most attractive candidates for this type of screening, would be prokaryotic organisms (because they grow quickly, are relatively simple to manipulate and because large numbers of clones can be created) which express and display at their surface a functional binding domain eg. an antibody, receptor, enzyme etc. In the UK patent GB 2137631B methods for the co-expression in a single host cell of the variable H and L chain genes of immunoglobulins were disclosed. However, the protein was expressed intracellularly and was insoluble. Further, the protein required extensive processing to generate antibody fragments with binding activity and this generated material with only a fraction of the binding activity expected for antibody fragments at this concentration. It has already been shown that antibody fragments can be secreted through bacterial membranes with the appropriate signal peptide (Skerra, A. and Pluckthun, A. 1988 Science 240 1038-1040; Better, M et al 1988, Science 240 1041-1043) with a consequent increase in the binding activity of antibody fragments. These methods require screening of individual clones for binding activity in the same way as do mouse monoclonal antibodies.

It has not been shown however, how a functional binding domain eg an antibody, antibody fragment, receptor, enzyme etc can be held on the bacterial surface in a configuration which allows sampling of say its antigen binding properties and selection for clones with desirable properties. In large part, this is because the bacterial surface is a complex structure, and in the gram-negative organisms there is an outer wall which further complicates the position. Further, it has not been shown that eg an antibody domain will fold correctly when expressed as a fusion with a surface protein of bacteria or bacteriophage.

Bacteriophage are attractive prokaryote related organisms for this type of screening. In general, their surface is a relatively simple structure, they can be grown easily in large numbers, they are amenable to the practical handling involved in many potential mass screening programmes, and they carry genetic information for their own synthesis within a small, simple package. The difficulty has been to practically solve the problem of how to use bacteriophages in this manner. A Genex Corporation patent application number W088/06630 has proposed that the bacteriophage lambda would be a suitable vehicle for the

expression of antibody molecules, but they do not provide a teaching which enables the general idea to be carried out. For example W088/06630 does not demonstrate that any sequences: (a) have been expressed as a fusion with gene V; (b) have been expressed on the surface of lambda; and (c) have been expressed so that the protein retains biological activity. Furthermore there is no teaching on how to screen for suitable fusions. Also, since the lambda virions are assembled within the cell, the fusion protein would be expressed intracellularly and would be predicted to be inactive. Bass et al., in December 1990 (after the earliest priority date for the present application) describe deleting part of gene III of the filamentous bacteriophage M13 and inserting the coding sequence for human growth hormone (hGH) into the N-terminal site of the gene. The growth hormone displayed by M13 was shown to be functional. (Bass, S., et al. Proteins, Structure, Function and Genetics (1990) 8: 309-314). A functional copy of gene III was always present in addition, when this fusion was expressed. A Protein Engineering Corporation patent application W090/02809 proposes the insertion of the coding sequence for bovine pancreatic trypsin inhibitor (BPTI) into gene VIII of M13. However, the proposal was not shown to be operative. For example, there is no demonstration of the expression of BPTI sequences as fusions with protein VIII and display on the surface of M13. Furthermore this document teaches that when a fusion is made with gene III, it is necessary to use a second synthetic copy of gene III, so that some unaltered gene III protein will be present. The embodiments of the present application do not do this. In embodiments where phagemid is rescued with M13K07 gene III deletion phage, there is no unaltered gene III present.

W090/02809 also teaches that phagemids that do not contain the full genome of M13 and require rescue by coinfection with helper phage are not suitable for these purposes because coinfection could lead to recombination.

In all embodiments where the present applicants have used phagemids, they have used a helper phage and the only sequences derived from filamentous bacteriophage in the phagemids are the origin of replication and gene III sequences.

W090/02809 also teaches that their process needed information such as nucleotide sequence of the starting molecule and its three-dimensioned structure. The use of a pre-existing repertoire of binding molecules to select for a binding member, such as is disclosed herein, for example using an immunoglobulin gene repertoire of animals, was not disclosed. Further, they do not discuss favouring variegation of their binding molecules in natural blocks of variation such as CDRs of immunoglobulins, in order to favour generation of improved molecules and prevent unfavourable variations. W090/02809 also specifically excluded the application of their process to the production of scFv molecules.

In each of the above discussed patents (W088/06630 and

WO90/02809), the protein proposed for display is a single polypeptide chain. There is no disclosure of a method for the display of a dimeric molecule by expression of one monomer as a fusion with a capsid protein and the other protein in a free form.

Another disclosure published in May 1991 (after the earliest priority date for the present application) describes the insertion into gene VIII of M13, the coding sequences for one of the two chains of the Fab portion of an antibody with co-expression of the other from a plasmid. The two chains were demonstrated as being expressed as a functional Fab fragment on the surface of the phage (Kang A.S. et al., (1991) Proc. Natl. Acad. Sci. USA, 88 p4363-4366). No disclosure was made of the site of insertion into gene VIII and the assay for pAb binding activity by ELISA used a reagent specific for antibody L chain rather than for phage. A further disclosure published in March 1991 (after the earliest priority date for the present application) describes the insertion of a fragment of the AIDS virus protein gag into the N-terminal portion of gene III of the bacteriophage fd. The expression of the gag protein fragment was detected by immunological methods, but it was not shown whether or not the protein was expressed in a functional form (Tsunetsugu-Yokota Y et al. (1991) Gene 99 p261-265).

The problem of how to use bacteriophages in this way is in fact a difficult one. The protein must be inserted into the phage in such a way that the integrity of the phage coat is not undermined, and the protein itself should be functional retaining its biological activity with respect to antigen binding. Thus, where the protein of choice is an antibody, it should fold efficiently and correctly and be presented for antigen binding. Solving the problem for antibody molecules and fragments would also provide a general method for any biomolecule which is a member of a specific binding pair e.g. receptor molecules and enzymes.

Surprisingly, the applicants have been able to construct a bacteriophage that expresses and displays at its surface a large biologically functional binding molecule (eg antibody fragments, and enzymes and receptors) and which remains intact and infectious. The applicants have called the structure which comprises a virus particle and a binding molecule displayed at the viral surface a 'package'. Where the binding molecule is an antibody, an antibody derivative or fragment, or a domain that is homologous to an immunoglobulin domain, the applicants call the package a 'phage antibody' (pAb). However, except where the context demands otherwise, where the term phage antibody is used generally, it should also be interpreted as referring to any package comprising a virus particle and a biologically functional binding molecule displayed at the viral surface.

pAbs have a range of applications in selecting antibody genes encoding antigen binding activities. For example, pAbs could be used for the cloning and rescue of hybridomas

(Orlandi, R., et al (1989) PNAS 86 p3833-3837), and in the screening of large combinatorial libraries (such as found in Huse, W.D. et al., 1989, Science 246, 1275-1281). In particular, rounds of selection using pAbs may help in rescuing the higher affinity antibodies from the latter libraries. It may be preferable to screen small libraries derived from antigen-selected cells (Casali, P., et al., (1986) Science 234 p476-479) to rescue the original VH/VL pairs comprising the Fv region of an antibody. The use of pAbs may also allow the construction of entirely synthetic antibodies. Furthermore, antibodies may be made which have some synthetic sequences e.g. CDRs, and some naturally derived sequences. For example, V-gene repertoires could be made in vitro by combining un-rearranged V genes, with D and J segments. Libraries of pAbs could then be selected by binding to antigen, hypermutated in vitro in the antigen-binding loops or V domain framework regions, and subjected to further rounds of selection and mutagenesis.

As previously discussed, separate H and L chain libraries lose the original pairing between the chains. It is difficult to make and screen a large enough library for a particularly advantageous combination of H and L chains.

For example, in a mouse there are approximately 10^7 possible H chains and 10^7 possible L chains. Therefore, there are 10^{14} possible combinations of H and L chains, and to test for anything like this number of combinations one would have to create and screen a library of about 10^{14} clones. This has not previously been a practical possibility.

The present invention provides a number of approaches which ameliorate this problem.

In a first approach, (a random combinatorial approach, see examples 20 and 21) as large a library as is practically possible is created which expresses as many of the 10^{14} potential combinations as possible. However, by virtue of the expression of the H and L chains on the surface of the phage, it is reasonably practicable to select the desired combination, from all the generated combinations by affinity techniques (see later for description of selection formats).

In a second approach (called a dual combinatorial approach by the present applicants, see example 26), a large library is created from two smaller libraries for selection of the desired combination. This ameliorates the problems still further. The approach involves the creation of: (i) a first library of say 10^7 e.g. H chains which are displayed on a bacteriophage (as a fusion with the protein encoded by gene III) which is resistant to e.g. tetracycline; and (ii) a second library of say 10^7 e.g. L chains in which the coding sequences for these light chains are within a plasmid vector containing an origin of replication for a bacteriophage (a phagemid) which is resistant to e.g. ampicillin (i.e. a different antibiotic) and are expressed in the periplasmic space of a host bacterium. The first library is then used to infect the bacteria containing the second library to provide 10^{14} combinations of H and L

chains on the surface of the resulting phage in the bacterial supernatant.

The advantage of this approach is that two separate libraries of eg 10^7 are created in order to produce 10^{14} combinations. Creating a 10^7 library is a practical possibility.

The 10^{14} combinations are then subjected to selection (see later for description of selection formats) as disclosed by the present application. This selection will then produce a population of phages displaying a particular combination of H and L chains having the desired specificity. The phages selected however, will only contain DNA encoding one partner of the paired H and L chains (deriving from either the phage or phagemid). The sample eluate containing the population is then divided into two portions. A first portion is grown on e.g. tetracycline plates to select those bacteriophage containing DNA encoding H chains which are involved in the desired antigen binding. A second portion is grown on e.g. ampicillin plates to select those bacteriophage containing phagemid DNA encoding L chains which are involved in the desired antigen binding. A set of colonies from individually isolated clones e.g. from the tetracycline plates are then used to infect specific colonies e.g. from the ampicillin plates. This results in bacteriophage expressing specific combinations of H and L chains which can then be assayed for antigen binding.

In a third approach (called a hierarchical dual combinational approach by the present applicants), an individual colony from either the H or L chain clone selected by growth on the antibiotic plates, is used to infect a complete library of clones encoding the other chain (H or L). Selection is as described above. This favours isolation of the most favourable combination.

In a fourth approach (called a hierarchical approach by the present applicants, see examples 22 and 46) both chains are cloned into the same vector. However, one of the chains which is already known to have desirable properties is kept fixed. A library of the complementary chain is inserted into the same vector. Suitable partners for the fixed chain are selected following display on the surface of bacteriophage.

In a fifth approach (see example 48), to improve the chances of recovering original pairs, the complexity of the combinatorial libraries can be reduced by using small B populations of B-lymphocytes selected for binding to a desired antigen. The cells provide e.g. mRNA or DNA, for preparing libraries of antibody genes for display on phage. This technique can be used in combination with the above mentioned four approaches for selection of antibody specificities.

Phagemids have been mentioned above. The applicants have realised and demonstrated that in many cases phagemids will be preferred to phage for cloning antibodies because it is easier to use them to generate more comprehensive

libraries of the immune repertoire. This is because the phagemid DNA is approximately 100 times more efficient than bacteriophage DNA in transforming bacteria (see example 19). Also, the use of phagemids gives the ability to vary the number of gene III binding molecule fusion proteins displayed on the surface of the bacteriophage (see example 17). For example, in a system comprising a bacterial cell containing a phagemid encoding a gene III fusion protein and infected with a helper phage, induction of expression of the gene III fusion protein to different extents, will determine the number of gene III fusion proteins present in the space defined between the inner and outer bacterial membranes following superinfection. This will determine the ratio of gene III fusion protein to native gene III protein displayed by the assembled phage.

Expressing a single fusion protein per virion may aid selection of antibody specificities on the basis of affinity by avoiding the 'avidity' effect where a phage expressing two copies of a low affinity antibody would have the same apparent affinity as a phage expressing one copy of a higher affinity antibody. In some cases however, it will be important to display all the gene III molecules derived by superinfection of cells containing phagemids to have fusions (e.g. for selecting low affinity binding molecules or improving sensitivity on ELISA). One way to do this is to superinfect with a bacteriophage which contains a defective gene III. The applicants have therefore developed and used a phage which is deleted in gene III. This is completely novel.

The demonstration that a functional antigen-binding domain can be displayed on the surface of phage, has implications beyond the construction of novel antibodies. For example, if other protein domains can be displayed at the surface of a phage, phage vectors could be used to clone and select genes by the binding properties of the displayed protein. Furthermore, variants of proteins, including epitope libraries built into the surface of the protein, could be made and readily selected for binding activities. In effect, other protein architectures might serve as "nouvelle" antibodies.

The technique provides the possibility of building antibodies from first principles, taking advantage of the structural framework on which the antigen binding loops fold. In general, these loops have a limited number of conformations which generate a variety of binding sites by alternative loop combinations and by diverse side chains. Recent successes in modelling antigen binding sites augurs well for de novo design. In any case, a high resolution structure of the antigen is needed. However, the approach is attractive for making e.g. catalytic antibodies, particularly for small substrates. Here side chains or binding sites for prosthetic groups might be introduced, not only to bind selectively to the transition state of the substrate, but also to participate directly in bond making and breaking. The only question is whether the antibody

architecture, specialised for binding, is the best starting point for building catalysts. Genuine enzyme architectures, such as the triose phosphate isomerase (TIM) barrel, might be more suitable. Like antibodies, TIM enzymes also have a framework structure (a barrel of β -strands and α -helices) and loops to bind substrate. Many enzymes with a diversity of catalytic properties are based on this architecture and the loops might be manipulated independently on the frameworks for design of new catalytic and binding properties. The phage selection system as provided by the present disclosure can be used to select for antigen binding activities and the CDR loops thus selected, used on either an antibody framework or a TIM barrel framework. Loops placed on a e.g. a TIM barrel framework could be further modified by mutagenesis and subjected to further selection. Thus, there is no need to select for high affinity binding activities in a single step. The strategy of the immune system, in which low affinity evolves to high affinity seems more realistic and can be mimicked using this invention.

One class of molecules that could be useful in this type of application are receptors. For example, a specific receptor could be displayed on the surface of the phage such that it would bind its ligand. The receptor could then be modified by, for example, in vitro mutagenesis and variants having higher binding affinity for the ligand selected. The selection may be carried out according to one or more of the formats described below with reference to figure 2 (which refers particularly to pAbs) in which the pAb antibody is replaced with a phage receptor and the antigen with a ligand 1.

Alternatively, the phage-receptor could be used as the basis of a rapid screening system for the binding of ligands, altered ligands, or potential drug candidates. The advantages of this system namely of simple cloning, convenient expression, standard reagents and easy handling makes the drug screening application particularly attractive. In the context of this discussion, receptor means a molecule that binds a specific, or group of specific, ligand(s). The natural receptor could be expressed on the surface of a population of cells, or it could be the extracellular domain of such a molecule (whether such a form exists naturally or not), or a soluble molecule performing a natural binding function in the plasma, or within a cell or organ.

Another possibility, is the display of an enzyme molecule or active site of an enzyme molecule on the surface of a phage (see examples 11,12,30,31,32 and 36). Once the phage enzyme is expressed, it can be selected by affinity chromatography, for instance on columns derivatized with transition state analogues. If an enzyme with a different or modified specificity is desired, it may be possible to mutate an enzyme displayed as a fusion on bacteriophage and then select on a column derivatised with an analogue selected to have a higher affinity for an enzyme with the desired modified specificity.

Although throughout this application, the applicants discuss the possibility of screening for higher affinity variants of pAbs, they recognise that in some applications, for example low affinity chromatography (Ohlson, S. et al Anal. Biochem. 169, p204-208 (1988)), it may be desirable to isolate lower affinity variants.

Examples 21 and 23 show that the present invention provides a way of producing antibodies with low affinities (as seen in the primary immune response or in unimmunised animals). This is made possible by displaying multiple copies of the antibody on the phage surface in association with gene III protein. Thus, pAbs allow genes for these antibodies to be isolated and if necessary, mutated to provide improved antibodies.

pAbs also allow the selection of antibodies for improved stability. It has been noted for many antibodies, that yield and stability are improved when the antibodies are expressed at 30°C rather than 37°C. If pAbs are displayed at 37°C, only those which are stable will be available for affinity selection. When antibodies are to be used in vivo for therapeutic or diagnostic purposes, increased stability would extend the half-life of antibodies in circulation.

Although stability is important for all antibodies and antibody domains selected using phage, it is particularly important for the selection of Fv fragments which are formed by the non-covalent association of VH and VL fragments. Fv fragments have a tendency to dissociate and have a much reduced half-life in circulation compared to whole antibodies. Fv fragments are displayed on the surface of phage, by the association of one chain expressed as a gene III protein fusion with the complementary chain expressed as a soluble fragment. If pairs of chains have a high tendency to dissociate, they will be much less likely to be selected as pAbs. Therefore, the population will be enriched for pairs which do associate stably. Although dissociation is less of a problem with Fab fragments, selection would also occur for Fab fragments which associate stably. pAbs allow selection for stability to protease attack, only those pAbs that are not cleaved by proteases will be capable of binding their ligand and therefore populations of phage will be enriched for those displaying stable antibody domains.

The technique of displaying binding molecules on the phage surface can also be used as a primary cloning system. For example, a cDNA library can be constructed and inserted into the bacteriophage and this phage library screened for the ability to bind a ligand. The ligand/binding molecule combination could include any pair of molecules with an ability to specifically bind to

one another e.g. receptor/ligand, enzyme/substrate (or analogue), nucleic acid binding protein/nucleic acid etc. If one member of the complementary pair is available, this may be a preferred way of isolating a clone for the other member of the pair.

It will often be necessary to increase the diversity of a population of genes cloned for the display of their proteins on phage or to mutate an individual nucleotide sequence. Although in vitro or in vivo mutagenesis techniques could be used for either purpose, a particularly suitable method would be to use mutator strains. A mutator strain is a strain which contains a genetic defect which causes DNA replicated within it to be mutated with respect to its parent DNA. Hence if a population of genes as gene III fusions is introduced into these strains it will be further diversified and can then be transferred to a non-mutator strain, if desired, for display and selection. Example 38 covers the use of mutator strains with phage antibodies (an example of in vitro mutagenesis and selection of phage antibodies is given in example 45).

Targeted gene transfer

A useful and novel set of applications makes use of the binding protein on the phage to target the phage genome to a particular cell or group of cells. For example, a pAb specific for a cell surface molecule could be used to bind to the target cell via the surface molecule. The phage could then be internalised, either through the action of the receptor itself or as the result of another event (e.g. an electrical discharge such as in the technique of electroporation). The phage genome would then be expressed if the relevant control signals (for transcription and translation and possibly replication) were present. This would be particularly useful if the phage genome contained a sequence whose expression was desired in the target cell (along with the appropriate expression control sequences). A useful sequence might confer antibiotic resistance to the recipient cell or label the cell by the expression of its product (e.g. if the sequence expressed a detectable gene product such as a luciferase, see White, M, et al, Techniques 2(4), p194-201 (1990)), or confer a particular property on the target cell (e.g. if the target cell was a tumour cell and the new sequence directed the expression of a tumour suppressing gene), or express an antisense construct designed to turn off a gene or set of genes in the target cell, or a gene or gene product designed to be toxic to the target cell.

Alternatively, the sequence whose expression is desired in the target cell can be encoded on a phagemid. The phagemid DNA may then be incorporated into a phage displaying an antibody specific for a cell surface

receptor. For example, incorporation may be by superinfection of bacteria containing the phagemid, with a helper phage whose genome encodes the antibody fragment specific for the target cell. The package is then used to direct the phagemid to the target cell.

This technique of "targeted gene transfer" has a number of uses in research and also in therapy and diagnostics. For example, gene therapy often aims to target the replacement gene to a specific cell type that is deficient in its activity. Targetting pAbs provide a means of achieving this.

In diagnostics, phage specific for particular bacteria or groups of bacteria have been used to target marker genes, e.g. luciferase, to the bacterial host (see, for example, Ulitzer, S., and Kuhn, J., EPA 85303913.9). If the host range of the phage is appropriate, only those bacteria that are being tested for, will be infected by the phage, express the luciferase gene and be detected by the light they emit. This system has been used to detect the presence of Salmonella. One major problem with this approach is the initial isolation of a bacteriophage with the correct host range and then the cloning of a luciferase gene cassette into that phage, such that it is functional. The pAb system allows the luciferase cassette to be cloned into a well characterised system (filamentous phage) and allows simple selection of an appropriate host range, by modifying the antibody (or other binding molecule) specificity that the pAb encodes.

The present applicants have also been able to develop novel selection systems and assay formats which depend on the unique properties of these replicable genetic display packages e.g. pAbs.

TERMINOLOGY

Much of the terminology discussed in this section has been mentioned in the text where appropriate.

Specific Binding Pair

This describes a pair of molecules (each being a member of a specific binding pair) which are naturally derived or synthetically produced. One of the pair of molecules, has an area on its surface, or a cavity which specifically binds to, and is therefore defined as complementary with a particular spatial and polar organisation of the other molecule, so that the pair have the property of binding specifically to each other. Examples of types of specific binding pairs are antigen-antibody, biotin-avidin, hormone-hormone receptor, receptor-ligand, enzyme-substrate, IgG-protein A.

Multimeric Member

This describes a first polypeptide which will associate with at least a second polypeptide, when the polypeptides are expressed in free form and/or on the

surface of a substrate. The substrate may be provided by a bacteriophage. Where there are two associated polypeptides, the associated polypeptide complex is a dimer, where there are three, a trimer etc. The dimer, trimer, multimer etc or the multimeric member may

comprise a member of a specific binding pair. Example multimeric members are heavy domains based on an immunoglobulin molecule, light domains based on an immunoglobulin molecule, T-cell receptor subunits.

Replicable Genetic Display Package (Rgdp)

This describes a biological particle which has genetic information providing the particle with the ability to replicate. The particle can display on its surface at least part of a polypeptide. The polypeptide can be encoded by genetic information native to the particle and/or artificially placed into the particle or an ancestor of it. The displayed polypeptide may be any member of a specific binding pair eg. heavy or light chain domains based on an immunoglobulin molecule, an enzyme or a receptor etc.

The particle may be a virus eg. a bacteriophage such as fd or M13.

Package

This describes a replicable genetic display package in which the particle is displaying a member of a specific binding pair at its surface. The package may be a bacteriophage which displays an antigen binding domain at its surface. This type of package has been called a phage antibody (pAb).

Antibody

This describes an immunoglobulin whether natural or partly or wholly synthetically produced. The term also covers any protein having a binding domain which is homologous to an immunoglobulin binding domain. These proteins can be derived from natural sources, or partly or wholly synthetically produced.

Example antibodies are the immunoglobulin isotypes and the Fab, $F(ab^1)_2$, scFv, Fv, dAb, Fd fragments.

Immunoglobulin Superfamily

This describes a family of polypeptides, the members of which have at least one domain with a structure related to that of the variable or constant domain of immunoglobulin molecules. The domain contains two β -sheets and usually a conserved disulphide bond (see A.F. Williams and A.N. Barclay 1988 Ann. Rev Immunol. 6 381-405).

Example members of an immunoglobulin superfamily are CD4, platelet derived growth factor receptor (PDGFR), intercellular adhesion molecule. (ICAM). Except where the context otherwise dictates, reference to immunoglobulins and immunoglobulin homologs in this application includes members of the immunoglobulin

superfamily and homologs thereof.

Homologs

5 This term indicates polypeptides having the same or conserved residues at a corresponding position in their primary, secondary or tertiary structure. The term also extends to two or more nucleotide sequences encoding the homologous polypeptides.

Example homologous peptides are the immunoglobulin isotypes.

10 Functional

In relation to a sbp member displayed on the surface of a rgdp, means that the sbp member is presented in a folded form in which its specific binding domain for its complementary sbp member is the same or closely analogous to its native configuration, whereby it exhibits similar specificity with respect to the complementary sbp member. In this respect, it differs from the peptides of Smith et al, supra, which do not have a definite folded configuration and can assume a variety of configurations determined by the complementary members with which they may be contacted.

Genetically diverse population

25 In connection with sbp members or polypeptide components thereof, this is referring not only to diversity that can exist in the natural population of cells or organisms, but also diversity that can be created by artificial mutation in vitro or in vivo.

30 Mutation in vitro may for example, involve random mutagenesis using oligonucleotides having random mutations of the sequence desired to be varied. In vivo mutagenesis may for example, use mutator strains of host microorganisms to harbour the DNA (see Example 38 below).

Domain

35 A domain is a part of a protein that is folded within itself and independently of other parts of the same protein and independently of a complementary binding member.

Folded Unit

40 This is a specific combination of an α -helix and/or β -strand and/or β -turn structure. Domains and folded units contain structures that bring together amino acids that are not adjacent in the primary structure.

Free Form

45 This describes the state of a polypeptide which is not displayed by a replicable genetic display package.

Conditionally Defective

50 This describes a gene which does not express a particular polypeptide under one set of conditions, but expresses it under another set of conditions. An example, is a gene containing an amber mutation expressed in non-suppressing or suppressing hosts respectively.

Alternatively, a gene may express a protein which is

defective under one set of conditions, but not under another set. An example is a gene with a temperature sensitive mutation.

Suppressible Translational Stop Codon

5 This describes a codon which allows the translation of nucleotide sequences downstream of the codon under one set of conditions, but under another set of conditions translation ends at the codon. Example of suppressible translational stop codons are the amber, ochre and opal
10 codons.

Mutator Strain

This is a host cell which has a genetic defect which causes DNA replicated within it to be mutated with respect to its parent DNA. Example mutator strains are
15 NR9046mutD5 and NR9046 mut T1 (see Example 38).

Helper Phage

This is a phage which is used to infect cells containing a defective phage genome and which functions to complement the defect. The defective phage genome can
20 be a phagemid or a phage with some function encoding gene sequences removed. Examples of helper phages are M13K07, M13K07 gene III no. 3; and phage displaying or encoding a binding molecule fused to a capsid protein.

Vector

25 This is a DNA molecule, capable of replication in a host organism, into which a gene is inserted to construct a recombinant DNA molecule.

Phage Vector

30 This is a vector derived by modification of a phage genome, containing an origin of replication for a bacteriophage, but not one for a plasmid.

Phagemid Vector

35 This is a vector derived by modification of a plasmid genome, containing an origin of replication for a bacteriophage as well as the plasmid origin of replication.

Secreted

40 This describes a rgdp or molecule that associates with the member of a sbp displayed on the rgdp, in which the sbp member and/or the molecule, have been folded and the package assembled externally to the cellular cytosol.

Repertoire of Rearranged Immunoglobulin Genes

45 A collection of naturally occurring nucleotides eg DNA sequences which encoded expressed immunoglobulin genes in an animal. The sequences are generated by the in vivo rearrangement of eg V, D and J segments for H chains and eg the V and J segments for L chains. Alternatively the sequences may be generated from a cell line immunised in vitro and in which the rearrangement in
50 response to immunisation occurs intracellularly.

Library

A collection of nucleotide eg DNA, sequences within

clones.

Repertoire of Artificially Rearranged Immunoglobulin Genes

5 A collection of nucleotide eg DNA, sequences derived wholly or partly from a source other than the rearranged immunoglobulin sequences from an animal. This may include for example, DNA sequences encoding VH domains by combining unrearranged V segments with D and J segments and DNA sequences encoding VL domains by combining V and
10 J segments.

Part or all of the DNA sequences may be derived by oligonucleotide synthesis.

Secretory Leader Peptide

15 This is a sequence of amino acids joined to the N-terminal end of a polypeptide and which directs movement of the polypeptide out of the cytosol.

Eluant

20 This is a solution used to breakdown the linkage between two molecules. The linkage can be a non-covalent or covalent bond(s). The two molecules can be members of a sbp.

Derivative

25 This is a substance which derived from a polypeptide which is encoded by the DNA within a selected rgdp. The derivative polypeptide may differ from the encoded polypeptide by the addition, deletion, substitution or insertion of amino acids, or by the linkage of other molecules to the encoded polypeptide. These changes may be made at the nucleotide or protein level. For example
30 the encoded polypeptide may be a Fab fragment which is then linked to an Fc tail from another source. Alternatively markers such as enzymes, flouresceins etc may be linked to eg Fab, scFv fragments.

35 The present invention provides a method for producing a replicable genetic display package or population such rgdps of which method comprises the steps of:

- 40 a) inserting a nucleotide sequence encoding a member of a specific binding pair eg. a binding molecule within a viral genome;
- b) culturing the virus containing said nucleotide sequence, so that said binding molecule is expressed and displayed by the virus at its surface.

45 The present invention also provides a method for selecting a rgdp specific for a particular epitope which comprises producing a population of such rgdps as described above and the additional step of selecting for said binding molecule by contac-ing the population with said epitope so that individual rgdps with the desired
50 specificity may bind to said epitope. The method may comprise one or more of the additional steps of: (i) separating any bound rgdps from the epitope; (ii)

recovering any separated rgdps and (iii) using the inserted nucleotide sequences from any separated rgdps in a recombinant system to produce the binding molecule separate from virus. The selection step may isolate the
 5 nucleotide sequence encoding the binding molecule of desired specificity, by virtue of said binding molecule being expressed in association with the surface of the virus in which said encoding nucleic acid is contained.

The present invention also provides a method of
 10 producing a multimeric member of a specific binding pair (sbp), which method comprises:
 expressing in a recombinant host organism a first polypeptide chain of said sbp member or a genetically diverse population of said sbp member fused to a
 15 component of a secreted replicable genetic display package (rgdp) which thereby displays said polypeptide at the surface of the package, and expressing in a recombinant host organism a second polypeptide chain of said multimer and causing or allowing the polypeptide
 20 chains come together to form said multimer as part of said rgdp at least one of said polypeptide chains being expressed from nucleic acid that is capable of being packaged using said component therefor, whereby the genetic material of each said rgdp encodes a said
 25 polypeptide chain.
 Both said chains may be expressed in the same host organism.

The first and second chains of said multimer may be expressed as separate chains from a single vector
 30 containing their respective nucleic acid.

At least one of said polypeptide chains may be expressed from a phage vector.

At least one of said polypeptide chains may be expressed from a phagemid vector, the method including
 35 using a helper phage, or a plasmid expressing complementing phage genes, to help package said phagemid genome, and said component of the rgdp is a capsid protein therefor. The capsid protein may be absent, defective or conditionally defective in the helper phage.

The method may comprise introducing a vector capable
 40 of expressing said first polypeptide chain, into a host organism which expresses said second polypeptide chain in free form, or introducing a vector capable of expressing said second polypeptide in free form into a host organism
 45 which expresses said first polypeptide chain.

Each of the polypeptide chain may be expressed from nucleic acid which is capable of being packaged as a rgdp using said component fusion product, whereby encoding
 50 nucleic acid for both said polypeptide chains are packaged in respective rgdps.

The nucleic acid encoding at least one of said first and second polypeptide chains may be obtained from a

library of nucleic acid including nucleic acid encoding said chain or a population of variants of said chain. Both the first and second polypeptide chains may be obtained from respective said libraries of nucleic acid.

5 The present invention also provides a method of producing a member of a specific binding pair (sbp), from a nucleic acid library including nucleic acid encoding said sbp member or a genetically diverse population of said type of sbp members, which method comprises:

10 expressing in recombinant host cells polypeptides encoded by said library nucleic acid fused to a component of a secreted replicable genetic display package (rgdp) or in free form for association with a polypeptide component of said sbp member which is
15 expressed as a fusion to said rgdp component so that the rgdp displays said sbp member in functional form at the surface of the package, said library nucleic acid being contained within the host cells in a form that is capable of being packaged using said rgdp
20 component, whereby the genetic material of an rgdp displaying an sbp member contains nucleic acid encoding said sbp member or a polypeptide component thereof.

25 The nucleotide sequences for the libraries may be derived from eg animal spleen cells or peripheral blood lymphocytes. Alternatively the nucleotide sequence may be derived by the in vitro mutagenesis of an existing antibody coding sequence.

30 The present invention also provides a method of producing a member of a specific binding pair (sbp), which method comprises:

35 expressing in recombinant host cells nucleic acid encoding said sbp member or a genetically diverse population of said type of sbp member wherein the or each said sbp member or a polypeptide component thereof is expressed as a fusion with a component of a secreted replicable genetic display package (rgdp) which displays said sbp member at the surface of the
40 package, nucleic acid encoding said sbp member or a polypeptide component thereof being contained within the host cell in a form that is capable of being packaged using said rgdp component whereby the genetic material of the rgdp displaying said sbp member encodes said sbp member or a polypeptide
45 component thereof, said host organism being a mutator strain which introduces genetic diversity into the sbp member to produce said mixed population.

50 The present invention also provides a method of producing a member of a specific binding pair (sbp), which method comprises:

expressing in recombinant host cells nucleic acid

encoding said sbp member or a genetically diverse population of said type of sbp member wherein the or each said sbp member or a polypeptide component thereof is expressed as a fusion with a component of a secreted replicable genetic display package (rgdp) which displays said sbp member in functional form at the surface of the package, nucleic acid encoding said sbp member or a polypeptide component thereof being contained within the host cell in a form that is capable of being packaged using said rgdp component whereby the genetic material of the rgdp displaying an sbp member encodes said sbp member or a polypeptide component thereof, said fusions being with bacteriophage capsid protein and the rgdps being formed with said fusions in the absence of said capsid expressed in wild-type form.

The present invention also provides a method of producing a member of a specific binding pair (sbp) which method comprises:

expressing in recombinant host cells nucleic acid encoding said sbp member or a genetically diverse population of said type of sbp member or a polypeptide component thereof fused to a component of a secreted replicable genetic display package (rgdp) which displays said sbp member in functional form at the surface of the package, nucleic acid encoding said sbp member or a polypeptide component thereof being contained within the host cell in a form that is capable of being packaged using said rgdp component whereby the genetic material of the rgdp displaying an sbp member or a polypeptide component thereof encodes said sbp member or a polypeptide component thereof, said sbp member or polypeptide component thereof being expressed from a phagemid as a capsid fusion, and a helper phage, or a plasmid expressing complementing phage genes, is used along with said capsid fusions to package the phagemid nucleic acid.

The library or genetically diverse population may be obtained from:

- (i) the repertoire of rearranged immunoglobulin genes of an animal immunised with complementary sbp member,
- (ii) the repertoire of rearranged immunoglobulin genes of an animal not immunised with complementary sbp member,
- (iii) a repertoire of artificially rearranged immunoglobulin gene or genes
- (iv) a repertoire of immunoglobulin homolog gene or genes; or
- (v) a mixture of any of (i), (ii), (iii) and (iv).

The capsid protein may be absent, defective or

conditionally defective in the helper phage.

The host cell may be a mutator strain which introduces genetic diversity into the sbp member nucleic acid.

5 The sbp member may comprise a domain which is, or is homologous to, an immunoglobulin domain.

The rgdp may be a bacteriophage, the host a bacterium, and said component of the rgdp a capsid protein for the bacteriophage. The phage may be a
10 filamentous phage. The phage may be selected from the class I phages fd, M13, fl, Ifl, lke, ZJ/Z, Ff and the class II phages Xf, Pfl and Pf3. The phage may be fd or a derivative of fd. The derivative may be tetracycline resistant. The said sbp member or polypeptide chain
15 thereof may be expressed as a fusion with the gene III capsid protein of phage fd or its counterpart in another filamentous phage. The sbp member or polypeptide chain thereof may be inserted in the N-terminal region of the mature capsid protein downstream of a secretory leader
20 peptide. The sequence may be inserted after amino acid +1 of the mature protein. The site for insertion may be flanked by short sequences corresponding to sequences which occur at each end of the nucleic acid to be inserted. For example where 4 the protein domain is an
25 immunoglobulin domain, the insertion site in the phage may be flanked by nucleotide sequences which code for the first five amino acids and the last five amino acids of the Ig domain. Such flanking nucleotide sequences are shown in figure 4(2) B and C, wherein the site-flanking
30 nucleotide sequences encode amino acid sequences QVQLQ and VTVSS which occur at either end of the VH domain, or QVQLQ and LEIKR which occur at either end of the Fv (combined VH + VL) domain. Each of these sequences flanking the insertion site may include a suitable
35 cleavage site, as shown in Fig 4.

Alternatively, the flanking nucleotide sequences shown in figure 4(2)B and C as described above, may be used to flank the insertion site for any nucleic acid to be inserted, whether or not that nucleic acid codes an
40 immunoglobulin.

The host may be E.coli.

Nucleic acid encoding an sbp member polypeptide may be linked downstream to a viral capsid protein through a suppressible translational stop codon.

45 As previously mentioned, the present invention also provides novel selection systems and assay formats. In these systems and formats, the gene sequence encoding the binding molecule (eg. the antibody) of desired specificity is separated from a general population of
50 rgdps having a range of specificities, by the fact of its binding to a specific target (eg the antigen or epitope). Thus the rgdps formed by said expression may be selected

or screened to provide an individual sbp member or a selected mixed population of said sbp members associated in their respective rgdps with nucleic acid encoding said sbp member or a polypeptide chain thereof. The rgdps may be selected by affinity with a member complementary to said sbp member.

Any rgdps bound to said second member may be recovered by washing with an eluant. The washing conditions may be varied in order to obtain rgdps with different binding affinities for said epitope. Alternatively, to obtain eg high affinity rgdps, the complementary member (eg an epitope) may be presented to the population of rgdps (eg pAbs) already bound to a binding member in which case pAbs with a higher affinity for the epitope will displace the already bound binding member. Thus the eluant may contain a molecule which competes with said rgdp for binding to the complementary sbp member. The rgdp may be applied to said complementary sbp member in the presence of a molecule which competes with said package for binding to said complementary sbp member. Nucleic acid derived from a selected or screened rgdp may be used to express said sbp member or a fragment or derivative thereof in a recombinant host organism. Nucleic acid from one or more rgdps may be taken and used to provide encoding nucleic acid in a further said method to obtain an individual sbp member or a mixed population of sbp members, or encoding nucleic acid therefor. The expression end product may be modified to produce a derivative thereof.

The expression end product or derivative thereof may be used to prepare a therapeutic or prophylactic medicament or a diagnostic product.

The present invention also provides recombinant host cells harbouring a library of nucleic acid fragments comprising fragments encoding a genetically diverse population of a type of member of a specific binding pair (sbp), each sbp member or a polypeptide component thereof being expressed as a fusion with a component of a secretable replicable genetic display package (rgdp), so that said sbp members are displayed on the surface of the rgdps in functional form and the genetic material of the rgdps encode the associated sbp member or a polypeptide component thereof. The type of sbp members may be immunoglobulins or immunoglobulin homologs, a first polypeptide chain of which is expressed as a said fusion with a component of the rgdp and a second polypeptide chain of which is expressed in free form and associates with the fused first polypeptide chain in the rgdp.

The present invention also provides a helper phage whose genome lacks nucleic acid encoding one of its capsid proteins, or whose encoding nucleic acid therefor is conditionally defective, or which encodes said capsid

protein in defective or conditionally defective form.

The present invention also provides a bacterial host cell containing a filamentous phage genome defective for a capsid protein thereof and wherein the host cell is capable of expressing capsid protein complementing said defect such that infectious phage particles can be obtained therefrom. The complementing capsid protein may be expressed in said host from another vector contained therein. The defective capsid protein may be gene III of phage fd or its counterpart in another filamentous phage.

The present invention also provides recombinant E.coli TG1 M13KO7 gIII No. 3 (NCTC 12478).

The present invention also provides a phage antibody having the form of a replicable genetic display package displaying on its surface in functional form a member of a specific binding pair or a specific binding domain thereof.

In the above methods, the binding molecule may be an antibody, or a domain that is homologous to an immunoglobulin. The antibody and/or domain may be either naturally derived or synthetic or a combination of both. The domain may be a Fab, scFv, Fv dAb or Fd molecule. Alternatively, the binding molecule may be an enzyme or receptor or fragment, derivative or analogue of any such enzyme or receptor. Alternatively, the binding molecule may be a member of an immunoglobulin superfamily and which has a structural form based on an immunoglobulin molecule.

The present invention also provides rgdps as defined above and members of specific binding pairs eg. binding molecules such as antibodies, enzymes, receptors, fragments and derivatives thereof, obtainable by use of any of the above defined methods. The derivatives may comprise members of the specific binding pairs fused to another molecule such as an enzyme or a Fc tail.

The invention also includes kits for carrying out the methods hereof. The kits will include the necessary vectors. One such vector will typically have an origin of replication for single stranded bacteriophage and either contain the sbp member nucleic acid or have a restriction site for its insertion in the 5' end region of the mature coding sequence of a phage capsid protein, and with a secretory leader coding sequence upstream of said site which directs a fusion of the capsid protein exogenous polypeptide to the periplasmic space.

The restriction sites in the vectors are preferably those of enzymes which cut only rarely in protein coding sequences.

The kit preferably includes a phagemid vector which may have the above characteristics, or may contain, or have a site for insertion, of sbp member nucleic acid for expression of the encoded polypeptide in free form.

The kits will also contain ancillary components required for carrying out the method, the nature of such components depending of course on the particular method employed.

5 Useful ancillary components may comprise helper phage, PCR primers, and buffers and enzymes of various kinds.

10 PCR primers and associated reagents for use where the sbp members are antibodies may have the following characteristics:

- (i) primers having homology to the 5' end of the sense or anti-sense strand of sequences encoding domains of antibodies; and
- 15 (ii) primers including tag sequences 5' to these homologous sequences which incorporate restriction sites to allow insertion into vectors; together with sequences to allow assembly of amplified VH and VL regions to enable expression as Fv, scFv or Fab fragments.

20 Buffers and enzymes are typically used to enable preparation of nucleotide sequences encoding Fv, scFv or Fab fragments derived from rearranged or unrearranged immunoglobulin genes according to the strategies described herein.

25 The applicants have chosen the filamentous F-specific bacteriophages as an example of the type of phage which could provide a vehicle for the display of binding molecules e.g. antibodies and antibody fragments and derivatives thereof, on their surface and facilitate subsequent selection and manipulation.

30 The F-specific phages (e.g. fl, fd and M13) have evolved a method of propagation which does not kill the host cell and they are used commonly as vehicles for recombinant DNA (Kornberg, A., DNA Replication, W.H. Freeman and Co., San Francisco, 1980). The single stranded DNA genome (approximately 6.4 Kb) of fd is extruded through the bacterial membrane where it sequesters capsid sub-units, to produce mature virions. These virions are 6 nm in diameter, 1µm in length and 40 each contain approximately 2,800 molecules of the major coat protein encoded by viral gene VIII and four molecules of the adsorption molecule gene III protein (g3p) the latter is located at one end of the virion. The structure has been reviewed by Webster et al., 1978 45 in The Single Stranded DNA Phages, 557-569, Cold Spring Harbor Laboratory Press. The gene III product is involved in the binding of the phage to the bacterial F-pilus.

50 Although these phages do not kill their host during normal replication, disruption of some of their genes can lead to cell death (Kornberg, A., 1980 supra.) This places some restraint on their use. The applicants have

recognized that gene III of phage fd is an attractive possibility for the insertion of biologically active foreign sequences. There are however, other candidate sites including for example gene VIII and gene VI.

5 The protein itself is only a minor component of the phage coat and disruption of the gene does not lead to cell death (Smith, G. 1988, *Virology* 167: 156-165). Furthermore, it is possible to insert some foreign sequences (with no biological function) into various positions within this gene (Smith, G. 1985 *Science* 228: 1315-1317., Parmley, S.F. and Smith, G.P. *Gene*: 73 (1988) p. 305-318., and de la Cruz, V.F., et al., 1988, *J. Biol. Chem.*, 263: 4318-4322). Smith et al described the display of peptides on the outer surface of phage but they did not describe the display of protein domains. Peptides can adopt a range of structures which can be different when in free solution, than when bound to, for example, an antibody, or when forming part of a protein (Stanfield, R.I. et al., (1990) *Science* 248, p712-719).

20 Proteins in general have a well defined tertiary structure and perform their biological function only when adopting this structure. For example, the structure of the antibody D1.3 has been solved in the free form and when bound to antigen (Bhat, T.N. et al., (1990) *Nature* 347, p483-485). The gross structure of the protein is identical in each instance with only minor variations around the binding site for the antigen. Other proteins have more substantial conformation changes on binding of ligand, for instance the enzymes hexokinase and pyruvate dehydrogenase during their catalytic cycle, but they still retain their overall pattern of folding. This structural integrity is not confined to whole proteins, but is exhibited by protein domains. This leads to the concept of a folded unit which is part of a protein, often a domain, which has a well defined primary, secondary and tertiary structure and which retains the same overall folding pattern whether binding to a binding partner or not. The only gene sequence that Smith et al., described that was of sufficient size to encode a domain (a minimum of perhaps 50 amino acids) was a 335bp fragment of a β -galactosidase corresponding to nucleotides 861-1195 in the β -galactosidase gene sequence (Parmley, S. + Smith, G.P. 1988 *supra*. This would encode 112 amino acids of a much larger 380 amino acid domain.

45 Therefore, prior to the present application, no substantially complete domain or folded unit had been displayed on phage. In these cases, although the infectivity of the virion was disrupted, the inserted sequences could be detected on the phage surface by use of e.g. antibodies.

The protein encoded by gene III has several domains (Pratt, D., et al., 1969 *Virology* 39:42-53., Grant, R.A.,

et al., 1981, J. Biol. Chem. 256: 539-546 and Armstrong, J., et al., FEBS Lett. 135: 167-172 1981.) including: (i) a signal sequence that directs the protein to the cell membrane and which is then cleaved off; (ii) a domain that anchors the mature protein into the bacterial cell membrane (and also the phage coat); and (iii) a domain that specifically binds to the phage receptor, the F-pilus of the host bacterium. Short sequences derived from protein molecules have been inserted into two places within the mature molecule (Smith, G., 1985 supra., and Parmley, S.F. and Smith G.P., 1988 supra.). Namely, into an inter-domain region and also between amino acids 2 and 3 at the N-terminus. The insertion sites at the N-terminus were more successful in maintaining the structural integrity of the gene III protein and displaying the peptides on the surface of the phage. By use of antisera specific for the peptides, the peptides inserted into this position were shown to be on the surface of the phage. These authors were also able to purify the phage, using this property. However, the peptides expressed by the phage, did not possess measurable biological functions of their own.

Retaining the biological function of a molecule when it is expressed in a radically different context to its natural state is difficult. The demands on the structure of the molecule are heavy. In contrast, retaining the ability to be bound by specific antisera is a passive process which imposes far less rigorous demands on the structure of the molecule. For example, it is the rule rather than the exception that polyclonal antisera will recognise totally denatured, and biologically inactive, proteins on Western blots (see for example, Harlow, E. and Lane, D., Antibodies, a Laboratory Manual, Cold Spring Harbor Laboratory Press 1988). Therefore, the insertion of peptides into a region that allows their structure to be probed with antisera teaches only that the region allows the inserted sequences to be exposed and does not teach that the region is suitable for the insertion of large sequences with demanding structural constraints for the display of a molecule with a biological or binding function. In particular, it does not teach that domains or folded units of proteins can be displayed from sequences inserted in this region.

This experience with Western blots is a graphic practical demonstration which shows that retaining the ability to be bound by specific antisera imposes far less rigorous demands on the structure of a polypeptide, than does folding for the retention of a biological function.

Studies have been carried out, in which E.coli have been manipulated to express the protein β -adrenergic receptor as a fusion with the outer membrane protein lamB. The β -adrenergic receptor was expressed in a

functional form as determined by the presence of binding activity. However, when an equivalent antibody fusion was made with lamB, the antibody fusion was toxic to the host cell.

3 The applicants have investigated the possibility of
 inserting the gene coding sequence for biologically
 active antibody fragments into the gene III region of fd
 to express a large fusion protein. As is apparent from
 10 the previous discussion, this approach makes onerous
 demands on the functionality of the fusion protein. The
 insertion is large, encoding antibody fragments of at
 least 100-200 amino acids; the antibody derived domain
 must fold efficiently and correctly to display antigen-
 15 binding; and most of the functions of gene III must be
 retained. The applicants approach to the construction of
 the fusion molecule was designed to minimise the risk of
 disrupting these functions. In an embodiment of the
 invention, the initial vector used was fd-tet (Zacher,
 A.N., et al., 1980, Gene 9, 127-140) a tetracycline
 20 resistant version of fd bacteriophage that can be
 propagated as a plasmid that confers tetracycline
 resistance to the infected *E.coli* host. The applicants
 chose to insert after the signal sequence of the fd gene
 III protein for several reasons. In particular, the
 25 applicants chose to insert after amino acid 1 of the
 mature protein to retain the context for the signal
 peptidase cleavage. To retain the structure and function
 of gene III itself, the majority of the original amino
 acids are synthesized after the inserted immunoglobulin
 30 sequences. The inserted immunoglobulin sequences were
 designed to include residues from the switch region that
 links VH-VL to CH1-CL (Lesk, A., and Chothia, C., Nature
 335, 188-190, 1988).

Surprisingly, by manipulating gene III of
 35 bacteriophage fd, the present applicants have been able
 to construct a bacteriophage that displays on its surface
 large biologically functional antibody, enzyme, and
 receptor molecules whilst remaining intact and
 infectious. Furthermore, the phages bearing antibodies
 40 of desired specificity, can be selected from a background
 of phages not showing this specificity.

The sequences coding for a population of antibody
 molecules and for insertion into the vector to give
 expression of antibody binding functions on the phage
 45 surface can be derived from a variety of sources. For
 example, immunised or non-immunised rodents or humans,
 and from organs such as spleen and peripheral blood
 lymphocytes. The coding sequences are derived from these
 sources by techniques familiar to those skilled in the
 50 art (Orlandi, R., et al., 1989 supra; Larrick, J.W., et
 al., 1989 supra; Chiang, Y.L., et al., 1989 Bio
 Techniques 7, p. 360-366; Ward, E.S., et al., 1989 supra;

Sastry, L., et al., 1989 supra.) or by novel linkage strategies described in examples 14, 33, 40 and 42. Novel strategies are described in examples 7, 25, 33, 39 and 40 for displaying dimeric molecules eg Fab and Fv fragments on the surface of a phage. Each individual pAb in the resulting library of pAbs will express antibodies or antibody derived fragments that are monoclonal with respect to their antigen-binding characteristics.

The disclosure made by the present applicants is important and provides a significant breakthrough in the technology relating to the production of biological binding molecules, their fragments and derivatives by the use of recombinant methods.

In standard recombinant techniques for the production of antibodies, an expression vector containing sequences coding for the antibody polypeptide chains is used to transform e.g. E.coli. The antibody polypeptides are expressed and detected by use of standard screening systems. When the screen detects an antibody polypeptide of the desired specificity, one has to return to the particular transformed E.coli expressing the desired antibody polypeptide. Furthermore, the vector containing the coding sequence for the desired antibody polypeptide then has to be isolated for use from E.coli in further processing steps.

In the present invention however, the desired antibody polypeptide when expressed, is already packaged with its gene coding sequence. This means that when the an antibody polypeptide of desired specificity is selected, there is no need to return to the original culture for isolation of that sequence. Furthermore, in previous methods in standard recombinant techniques, each clone expressing antibody needs to be screened individually. The present application provides for the selection of clones expressing antibodies with desired properties and thus only requires screening of clones from an enriched pool.

Because a rgdp (eg a pAb) is a novel structure that displays a member of a specific binding pair (eg. an antibody of monoclonal antigen-binding specificity) at the surface of a relatively simple replicable structure also containing the genetic information encoding the member, rgdps eg pAbs, that bind to the complementary member of the specific binding pair (eg antigen) can be recovered very efficiently by either eluting off the complementary member using for example diethylamine, high salt etc and infecting suitable bacteria, or by denaturing the structure, and specifically amplifying the sequences encoding the member using PCR. That is, there is no necessity to refer back to the original bacterial clone that gave rise to the pAb.

For some purposes, for example immunoprecipitation

and some diagnostic tests, it is advantageous to use polyclonal antibodies or antibody fragments. The present invention allows this to be achieved by either selection of an enriched pool of pAbs with desired properties or by mixing individually isolated clones with desired properties. The antibodies or antibody fragments may then be expressed in soluble form if desired. Such a selected polyclonal pAb population can be grown from stocks of phage, bacteria containing phagemids or bacteria expressing soluble fragments derived from the selected polyclonal population. Thus a reagent equivalent to a polyclonal antiserum is created which can be replicated and routinely manufactured in culture without use of animals.

15 SELECTION FORMATS AND AFFINITY MATURATION

Individual rgdps eg pAbs expressing the desired specificity eg for an antigen, can be isolated from the complex library using the conventional screening techniques (e.g. as described in Harlow, E., and Lane, D., 1988, supra Gherardi, E et al. 1990. J. Immunol. meth. 126 p61-68).

The applicants have also devised a series of novel selection techniques that are practicable only because of the unique properties of rgdps. The general outline of some screening procedures is illustrated in figure 2 using pAbs as an example type of rgdp.

The population/library of pAbs to be screened could be generated from immunised or other animals; or be created in vitro by mutagenising pre-existing phage antibodies (using techniques well-known in the art such as oligonucleotide directed mutagenesis (Sambrook, J., et al., 1989 Molecular Cloning a Laboratory Manual, Cold Spring Harbor Laboratory Press). This population can be screened in one or more of the formats described below with reference to figure 2, to derive those individual pAbs whose antigen binding properties are different from sample c.

Binding Elution

Figure 2(i) shows antigen (ag) bound to a solid surface (s) the solid surface (s) may be provided by a petri dish, chromatography beads, magnetic beads and the like. The population/library of pAbs is then passed over the ag, and those individuals p that bind are retained after washing, and optionally detected with detection system d. A detection system based upon anti-fd antisera is illustrated in more detail below in example 4. If samples of bound population p are removed under increasingly stringent conditions, the binding affinity represented in each sample will increase. Conditions of increased stringency can be obtained, for example, by increasing the time of soaking or changing the pH of the soak solution, etc.

Competition

Referring to figure 2(ii) antigen ag can be bound to a solid support s and bound to saturation by the original binding molecule c. If a population of mutant pAb (or a set of unrelated pAbs) is offered to the complex, only those that have higher affinity for antigen ag than c will bind. In most examples, only a minority of population c will be displaced by individuals from population p. If c is a traditional antibody molecule, all bound material can be recovered and bound p recovered by infecting suitable bacteria and/or by use of standard techniques such as PCR.

An advantageous application is where ag is used as a receptor and c the corresponding ligand. The recovered bound population p is then related structurally to the receptor binding site/and or ligand. This type of specificity is known to be very useful in the pharmaceutical industry.

Another advantageous application is where ag is an antibody and c its antigen. The recovered bound population p is then an anti-idiotypic antibody which have numerous uses in research and the diagnostic and pharmaceutical industries.

At present it is difficult to select directly for anti-idiotypic antibodies. pAbs would give the ability to do this directly by binding pAb libraries (eg a naive library) to B cells (which express antibodies on their surface) and isolating those phage that bound well.

In some instances it may prove advantageous to pre-select population p. For example, in the anti-idiotypic example above, p can be absorbed against a related antibody that does not bind the antigen.

However, if c is a pAb, then either or both c and p can advantageously be marked in some way to both distinguish and select for bound p over bound c. This marking can be physical, for example, by pre-labelling p with biotin; or more advantageously, genetic. For example, c can be marked with an EcoB restriction site, whilst p can be marked with an EcoK restriction site (see Carter, P. et al., 1985, Nucl. Acids Res. 13, 4431-4443). When bound p+c are eluted from the antigen and used to infect suitable bacteria, there is restriction (and thus no growth) of population c (i.e. EcoB restricting bacteria in this example). Any phage that grew, would be greatly enriched for those individuals from p with higher binding affinities. Alternatively, the genetic marking can be achieved by marking p with new sequences, which can be used to specifically amplify p from the mixture using PCR.

Since the bound pAbs can be amplified using for example PCR or bacterial infection, it is also possible to rescue the desired specificity even when insufficient

individuals are bound to allow detection via conventional techniques.

The preferred method for selection of a phage displaying a protein molecule with a desired specificity or affinity will often be elution from an affinity matrix with a ligand (eg example 21). Elution with increasing concentrations of ligand should elute phage displaying binding molecules of increasing affinity. However, when eg a pAb binds to its antigen with high affinity or avidity (or another protein to its binding partner) it may not be possible to elute the pAb from an affinity matrix with molecule related to the antigen. Alternatively, there may be no suitable specific eluting molecule that can be prepared in sufficiently high concentration. In these cases it is necessary to use an elution method which is not specific to eg the antigen-antibody complex. Some of the non-specific elution methods generally used reduce phage viability for instance, phage viability is reduced with time at pH12 (Rossomando, E.F. and Zinder N.D. J. Mol.Biol. 36 387-399 1968). There may be interactions between eg antibodies and affinity matrices which cannot be disrupted without completely removing phage infectivity. In these cases a method is required to elute phage which does not rely on disruption of eg the antibody - antigen interaction. A method was therefore devised which allows elution of bound pAbs under mild conditions (reduction of a dithiol group with dithiothreitol) which do not disrupt phage structure (example 47).

This elution procedure is just one example of an elution procedure under mild conditions. A particularly advantageous method would be to introduce a nucleotide sequence encoding amino acids constituting a recognition site for cleavage by a highly specific protease between the foreign gene inserted, in this instance a gene for an antibody fragment, and the sequence of the remainder of gene III. Examples of such highly specific proteases are Factor X and thrombin. After binding of the phage to an affinity matrix and elution to remove non-specific binding phage and weak binding phage, the strongly bound phage would be removed by washing the column with protease under conditions suitable for digestion at the cleavage site. This would cleave the antibody fragment from the phage particle eluting the phage. These phage would be expected to be infective, since the only protease site should be the one specifically introduced. Strongly binding phage could then be recovered by infecting eg. E.coli TG1 cells.

An alternative procedure to the above is to take the affinity matrix which has retained the strongly bound pAb and extract the DNA, for example by boiling in SDS solution. Extracted DNA can then be used to directly

transform E.coli host cells or alternatively the antibody encoding sequences can be amplified, for example using PCR with suitable primers such as those disclosed herein, and then inserted into a vector for expression as a soluble antibody for further study or a pAb for further rounds of selection.

Another preferred method for selection according to affinity would be by binding to an affinity matrix containing low amounts of ligand.

If one wishes to select from a population of phages displaying a protein molecule with a high affinity for its ligand, a preferred strategy is to bind a population of phage to an affinity matrix which contains a low amount of ligand. There is competition between phage, displaying high affinity and low affinity proteins, for binding to the ligand on the matrix. Phage displaying high affinity protein is preferentially bound and low affinity protein is washed away. The high affinity protein is then recovered by elution with the ligand or by other procedures which elute the phage from the affinity matrix (example 35 demonstrates this procedure).

In summary then, for recovery of the packaged DNA from the affinity step, the package can be simply eluted, it can be eluted in the presence of a homologous sbp member which competes with said package for binding to a complementary sbp member; it could be removed by boiling, it could be removed by proteolytic cleavage of the protein; and other methods will be apparent to those skilled in the art eg. destroying the link between the substrate and complementary sbp member to release said packaged DNA and sbp member. At any rate, the objective is to obtain the DNA from the package so that it can be used directly or indirectly, to express the sbp member encoded thereby.

The efficiency of this selection procedure for pAbs and the ability to create very large libraries means that the immunisation techniques developed to increase the proportion of screened cells producing antibodies of interest will not be an absolute requirement. The technique allows the rapid isolation of binding specificities eg antigen-binding specificities, including those that would be difficult or even unobtainable by conventional techniques, for example, catalytic or anti-idiotypic antibodies. Removal of the animal altogether is now possible, once a complete library of the immune repertoire has been constructed.

The novel structure of the pAb molecule can be used in a number of other applications, some examples of which are:
Signal Amplification

Acting as a novel molecular entity in itself, rgdps eg pAbs combine the ability to bind a specific molecule eg antigen with amplification, if the major coat protein

is used to attach another moiety. This moiety can be attached via immunological, chemical, or any other means and can be used, for example, to label the complex with detection reagents or cytotoxic molecules for use in vivo or in vitro.

Physical Detection

The size of the rgdps eg pAbs can be used as a marker particularly with respect to physical methods of detection such as electron microscopy and/or some biosensors, e.g. surface plasmon resonance.

Diagnostic Assays

The rgdps eg pAbs also have advantageous uses in diagnostic assays, particularly where separation can be effected using their physical properties for example centrifugation, filtration etc.

In order that the invention is more fully understood, embodiments will now be described in more detail by way of example only and not by way of limitation with reference to the figures described below.

Figure 1 shows the basic structure of the simplest antibody molecule IgG.

Figure 2 shows schematically selection techniques which utilise the unique properties of pAbs; 2i) shows a binding/elution system; and (2ii) shows a competition system (p=pAb; ag=antigen to which binding by pAb is required; c=competitor population e.g. antibody, pAb, ligand; s=substrate (e.g. plastic beads etc); d=detection system.

Figure 3 shows the vector fd-tet and a scheme for the construction of vectors, fdTPs/Bs (for insertion of VH coding sequences) and fdTPs/Xh for the insertion of scFv coding sequences.

Figure 4 shows the nucleotide sequences for the oligonucleotides and vectors. All sequences are drawn 5' to 3' and are numbered according to Beck et al., 1978, Nucl. Acid Res., 5: 4495-4503. 4.1 shows the sequences of the oligonucleotides used for mutagenesis (oligo's 1 and 2) or sequencing (oligo 3). The sequences shown were synthesized on an Applied Biosystems, oligonucleotide synthesizer and are complementary to the single stranded form of fd-tet (they are in the anti-sense form with respect to gene III). 4.2 shows the sequences of the various constructs around the gene III insertion site. These sequences are drawn in the sense orientation with respect to gene III; (A) fd-tet (and fdT6Bst) (B) fdTPs/Bs and (C) fdTPs/Xh. The key restriction enzyme sites are shown along with the immunoglobulin amino acids contributed by the vectors, (amino acid single letter code is used, see Harlow, E., and Lane, D., 1988 supra.).

Figure 5 shows the nucleotide and amino acid sequences for scFv in the vector scFvD1.3 myc. This gives the sequence of the anti-lysozyme single chain Fv

and surrounding sequences in scFvD1.3 myc, showing the N-terminal pel B signal peptide sequence and the C-terminal myc tag sequence (Ward, E.S., et al., 1989, supra.). Also shown is the peptide sequence linking the VH and VL regions. The amino acid sequence is represented above the nucleotide sequence by the single letter code, see Harlow, E., and Lane D., 1988 supra.

Figure 6 shows the binding of pAbs to lysozyme and the effect of varying the amount of supernatant. Each point is the average of duplicate samples. Lysozyme was coated at 1 mg/ml in 50 mM NaHCO₃.

Figure 7 shows the effect of varying the coating concentration of lysozyme or bovine serum albumin on the binding of pAbs to lysozyme in graphical form. Each point is the average of duplicate samples.

Figure 8 shows the sequence around the cloning site in gene III of fd-CAT2. Restriction enzyme sites are shown as well as the amino acids encoded by antibody derived sequences. These are flanked at the 5' end by the gene III signal peptide and at the 3' end by 3 alanine residues (encoded by the Not I restriction site) and the remainder of the mature gene III protein. The arrow shows the cleavage site for cutting of the signal peptide.

Figure 9 shows the binding of pAb (1.3) to lysozymes. Binding of phage as detected by ELISA to (a) hen egg-white lysozyme (HEL) (b) turkey egg-white lysozyme (TEL), (c) human lysozyme (HUL), (d) bovine serum albumin (BSA). A further control of (e) fdTPs/Bs to HEL.

Figure 10 shows a map of FabD1.3 in pUC19.

Figure 11 shows the ELISA results providing a comparison of lysozyme-binding by phage-Fab and phage-scFv. Vector=fdCAT2 (example 5); fdscFv(OX)=pAbNQ11 (Example 9); fdVHCH1 (D1.3)=grown in normal cells (i.e. no L chain, see example 7); fdFab(D1.3) i.e. fdVHCH1 (D1.3) grown in cells containing D1.3 L chain; fdscFv (D1.3)=pAbD1.3.

Figure 12 shows oligonucleotide probing of affinity purified phage. 10^{12} phage in the ratio of 1 pAb (D1.3) in 4×10^4 fdTPS/Bs phages were affinity purified and probed with an oligonucleotide specific for pAb (D1.3) A is a filter after one round of affinity purification (900 colonies total) and B is a filter after two rounds (372 colonies total).

Figure 13 shows the sequence of the anti-oxazolone antibody fragment NQ11 scFv. The sequence contributed by the linker is shown in the lower case. The sequence for VH is before the linker sequence and the sequence for VL is after the linker.

Figure 14 shows the ELISA results for binding pAb NQ11 and pAb D1.3 and vector fdTPs/xh to specified

antigens.

Figure 15 shows the sequence surrounding the phoA insertion in fd-phoAla166. The restriction sites used for cloning are shown, as well as the amino acids encoded by phoA around the insertion site. The first five amino acids of the mature fusion come from gene III.

Figure 16(1) shows the structure of gene III and the native BamHI site into which a scFv coding sequence was inserted in example 13 and figure 16(2) shows the natural peptide linker sites A and B for possible insertion of scFv coding sequences.

Figure 17 shows schematically the protocol for PCR assembly of mouse VH and VLK repertoires for phage display described in example 14.

Figure 18 shows examples of the final products obtained with the procedure of example 14. Lanes a and b show the products of the initial PCR using heavy and light chain primers respectively; lane c shows the complete assembled 700bp product before final digestion with NotI and ApaLI; M1, M2 markers ϕ 174 Hae III digest and 123 base pair ladder (BRL Limited, P.O. Box 35, Washington Road, Paisley, Scotland) respectively.

Figure 19 shows the binding of 125 I-PDGF-BB to fd h-PDGFB-R phage in immunoprecipitation assay and comparison to fdTPs/BS and no phage controls; binding is expressed as a percentage of the total 125 I-PDGF-BB added to the incubation.

Figure 20 shows the displacement of 125 I-PDGF-BB bound to fd-h-PDGFB-R phage using unlabelled PDGF-BB measured using an immunoprecipitation assay. Binding is expressed as a percentage of the total 125 I-PDGF-BB added to the incubation.

Figure 21 shows the displacement of 125 I-PDGF-BB bound to fd-h-PDGFB-R phage using unlabelled PDGF-BB measured using an immunoprecipitation assay. Non-specific binding of 125 I-PDGF-BB to vector phage fdTPs/BS in the absence of added unlabelled PDGF was deducted from each point.

Figure 22 shows the results of an ELISA of lysozyme binding by pCAT-3 scFv D1.3 phagemid in comparison with pCAT-3 vector (both rescued by M13K07) and fdCAT2 scFv D1.3 as described in example 17. The ELISA was performed as described in example 6 with modifications detailed in example 18.

Figure 23 shows the digestion pattern seen when individual clones, selected at random from a library of single chain Fv antibody genes derived from an immunised mouse; are digested with BstNI.

Figure 24 shows VH and VK gene sequences derived from the combinatorial library in example 21 and the hierarchical library in example 22.

Figure 25 shows a matrix of ELISA signals for clones

derived from random combinatorial library. Designation of the clones is as in figure 24. The number of clones found with each combination is shown by the numerals.

Figure 26 shows a) the phagemid pHEN1 a derivative of pUC119 described in example 24; and b) the cloning sites in the phagemid pHEN.

Figure 27. The antibody constructs cloned into fd-CAT2 and pHEN1 for display on the surface of phage. Constructs I, II, III and IV were cloned into both fd-CAT2 (as ApaLI-NotI fragments) and pHEN1 (as SfiI-NotI fragments) and pHEN1 (as SfiI-NotI fragments). All the constructs contained the heavy chain (VH) and light chain (VK) variable regions of the mouse anti-phOx antibody NQ10.12.5. The constant domains were human CK and CH1 (γ 1 isotype).

Figure 28. Three ways of displaying antibody fragments on the surface of phage by fusion to gene III protein.

Figure 29. Western blot of supernatant taken from pHEN1-II(+) or pHEN1(-) cultures in E.coli HB2151, showing secretion of Fab fragment from pHEN1-II only. The anti-human Fab detects both H and L chain. Due to the attached c-myc tag, the L chain, highlighted by both anti-c-myc tag and anti-human CK antisera, is slightly larger (calculated Mr 24625) than the H chain (calculated Mr23145).

Figure 30 is a plot showing the effect of lysozyme dilution on ratio of ELISA signals obtained using pAbD1.3 or soluble scFv D1.3.

Figure 31 is a plot showing the effect of lysozyme dilution on ELISA signals obtained using fdTscFvD1.3 and soluble scFvD1.3.

Figure 32 is a plot showing positive results from an ELISA screen of phage displaying scFv fragments derived from the cell line 013 which express a monoclonal antibody directed against oestriol.

Figure 33 is a plot showing positive results from an ELISA screen of phage displaying scFv fragments derived from the cell line 014 which express a monoclonal antibody directed against oestriol.

Figure 34 is a Western Blot showing expression of the alkaline phosphatase-gene 3 fusion. 16 μ l of 50 fold concentrate of each phage sample was detected on western blots with either anti-gene 3 antiserum (e-f) or with anti-alkaline phosphatase antiserum (c-f)

- a) fd-phoAla166 grown in TG1 cells
- b) fd-phoAla166 grown in KS272 cells
- c) fdCCAT2 grown in TG1 cells
- d) fdCAT2 grown in TG1 cells, mixed with 13 ng of purified alkaline phosphatase
- e) fd-phoAla166 grown in TG1 cells
- f) fdCAT2 grown in TG1 cells.

Figure 35 is a Western Blot showing ultrafiltration of phage-enzyme 100 μ l of 50 fold concentrate of phage (representing 5mls of culture supernatant) was centrifuged through ultrafiltration membranes with nominal molecular weight retention of 300,000 daltons. Western blots of flow through and retentate fractions were detected with anti-alkaline phosphatase antiserum. The equivalent of 800 μ l of original culture supernatant was run on the gel.

10 A. Phage were grown in TG1 cells. a) fd-phoAla166 before ultrafiltration (short exposure). b) fd-phoAla166 before ultrafiltration. c) fd-phoAla166 material retained on ultrafiltration membrane.
 15 B. Phage were grown in KS272 cells. a) fd-phoAla166 before ultrafiltration. b) fd-phoAla166 material retained on ultrafiltration membrane. c) fdCAT2. d) fdCAT2 mixed with purified alkaline phosphatase before ultrafiltration. e) Retentate from sample d. f) Flow through from sample d.

20 Figure 36 Electrophoresis of samples from stages of a Fab assembly. Samples from different stages in the PCR Fab assembly process described in example 33 were subjected to electrophoresis on a 1% TAE-agarose gel. Samples from a comparable scFv assembly process (as in
 25 example 14) are shown for comparison. Samples left to right are:

M	□	Markers
VHCH1	□	sequences encoding VHCH1 domains amplified by PCR
30 VKCK	□	sequences encoding VKCK domains amplified by PCR
-L	□	Fab assembly reaction performed in absence of linker
+L	□	Fab PCR assembly reaction product VHCH1 plus VKCK plus linker
35 M	□	Markers
VK	□	sequences encoding VK domain amplified by PCR
40 VL	□	sequences encoding VH domains amplified by PCR
-L	□	scFv assembly reaction in absence of linker
+L	□	scFv assembly reaction in presence of linker
45 M	□	Markers

Figure 37. Comparison of ELISA signals with scFv D1.3 cloned in fd-CAT2 (fd) or pCAT-3. pCAT-3 scFv1.3 has been rescued with M13K07 (K07). M13K07 Δ gIII No 3 (gIII No 3) or M13K07 gIIIANo 2 (gIIINo2). Phage antibodies are compared at 10 times (10x) 1 times (1x) or 0.1 times

(0.1x) concentrations relative to concentration in the supernatant after overnight growth. The fdCAT2 and pCAT-3 non-recombinant vector signals were <0.01 at 10x concentration. M13K07 gIIIΔNo 1 did not rescue at all, as judged by no signal above background in this ELISA.

Figure 38. Western blot of PEG precipitated phage used in ELISA probed with anti-g3p. Free g3p and the g3p-scFvD1.3 fusion bands are arrowed.

Sample 1 - fd scFvD1.3

Sample 2 - pCAT3 vector

Sample 3 - pCAT3 scFvD1.3 rescued with M13K07, no IPTG

Sample 4 - pCAT3 scFvD1.3 rescued with M13K07, 50μM IPTG

Sample 5 - pCAT3 scFvD1.3 rescued with M13K07, 100μM IPTG

Sample 6 - pCAT3 scFvD1.3 rescued with M13K07 gIIIΔNo3 (no IPTG)

Sample 7 - pCAT3 scFvD1.3 rescued with M13K07 gIIIΔNo 2 (no IPTG)

Panel A samples contain the equivalent of 8μl of phagemid culture supernatant per track, and 80μl of the fd supernatant (10-fold lower phage yield than the phagemid). Panel B phagemid samples are those used in panel A at a five-fold higher sample loading (equivalent to 40μl of culture supernatant per track) to enable visualisation of the fusion band in samples rescued with parental M13K07.

Figure 39 is a graph showing fdCAT2scFvD1.3 enrichment produced from a mixture of fdCAT2scFvD1.3 and fdCAT2TPB4 by one round of panning.

Figure 40 is a graph showing fdCAT2scFvD1.3 enrichment produced from a mixture of fdCAT2scFvD1.3 and fdCAT2TPB1 by one round of panning.

Figure 41. Western blot of phage proteins of fdCAT2(1) and fd-tet-SNase(2) with anti-g3p antiserum. Marker molecular weights bands are indicated(kD).

Figure 42. Nuclease assay of soluble SNase (3 ng)(A-1), fd-tet-SNase(4×10^9 TU, (B-1), fd-CAT2(2×10^{10} TU)(C-1) and of a PEG-precipitated fdCAT2 and SNase mixture(2×10^{10} TU and 0.7μg)(D-1) in a 10-fold dilution series (1 to 3 or 4). Marker (M) is a HindIII digest of λ-DNA(New England Biolabs).

Figure 43. ELISA signals obtained with fd-tet, fd-CD4-V1 and fd-CD4-V1V2. In each group of three, the samples are left to right phage concentrate(SN); phage concentrate plus soluble CD4(SN + sCD4); phage concentrate plus gp 120 (SN + gp 120).

Figure 44. shows the DNA sequence of scFv B18 (anti-NP).

Figure 45 shows a map of the insert of sequences encoding FvD1.3 present in fd-tet FvD1.3 (example 39). rbs designates the ribosome binding site. Gene III is now shown in its full length.

Figure 46. shows an ELISA assay of phages displaying

FvDl.3 or scFvDl.3 by binding to plates coated with lysogyme. Signals obtained at various dilution factors are shown. FvDl.3 (Δ S-Stuffer) which does not express Fv was used as a control.

5 Figure 47. shows a schematic representation of steps involved in the PCR assembly of nucleotide sequences encoding human Fab fragments. Details are in example 40.

10 Figure 48. shows A. a map of plasmid pJM1-FabDl.3 which is used for the expression of soluble human Fab fragments and as a template for the synthesis of linker DNA for Fab assembly. B. a schematic representation of sequences encoding a Fab construct. C. The sequence of DNA template for the synthesis of linker DNA for Fab assembly.

15 Figure 49. shows a schmatic representation of steps involved in the PCR assembly of nucleotide sequences encoding human scFv fragments. Details are in example 42.

20 Figure 50. ELISA assay of phage antibodies using plates coated with turkey egg lysogyme. Two clones B1 and A4 are shown derived by mutagenesis and selection from pAbDl.3 (example 45). Concentration (x axis) refers to the concentration of phage for each sample relative to the concentration in culture supernatant. B1 has raised binding to turkey egg lysogyme compared to Dl.3. A4 has reduced binding to hen egg lysogyme compared to Dl.3.

25 Figure 51. ELISA of phage antibodies binding to HEL and TEL. Clone 1 is fdCAT2scFvDl.3. Clones 2 to 10 were obtained from the library (example 46) after selection. The background values as defined by binding of these clones to BSA were subtracted.

30 Figure 52. shows the DNA sequence of the light chains Dl.3 M1F and M21 derived by selection from a hierarchical library in example 46.

35 Figure 53 shows a Fv lambda expression vector (example 48) derived from pUC119. It contains the rearranged lambda1 germ line gene. The heavy and light chain cassettes each contain a ribosome binding site upstream of the pel B leader (Restriction sites shown as: H=Hind III; Sp=SphI; B=BamHI, E=EcoRI).

40 Materials and Methods

The following procedures used by the present applicants are described in Sambrook, J. et al., 1989
45 supra.: restriction digestion, ligation, preparation of competent cells (Hanahan method), transformation, analysis of restriction enzyme digestion products on agarose gels, purification of DNA using phenol/chloroform, 5'-end labelling of oligonucleotides, filter screening of bacterial colonies, preparation of
50 2xTY medium and plates, preparation of tetracycline stock solutions, PAGE of proteins, preparation of phosphate buffered saline.

All enzymes were supplied by New England Biolabs (CP Laboratories, PO Box 22, Bishop's Stortford, Herts., England) and were used according to manufacturer's instructions unless otherwise stated.

The vector fd-tet (Zacher, A.N. et al., 1980, supra) was obtained from the American Type Culture Collection (ATCC No. 37000) and transformed into competent TG1 cells (genotype: K12 δ (lac-pro), sup E, thi, hsdD5/F traD36, pro A+B+, Lac I^q, lac δ M15).

Viral particles were prepared by growing TG1 cells containing the desired construct in 10 to 100 mls 2xTY medium with 15 μ g/ml tetracycline for 16-24 hours. The culture supernatant was collected by centrifugation for 10 mins at 10,000 rpm in an 8 x 50 ml rotor, Sorval RC-5B centrifuge. Phage particles were precipitated by adding 1/5th volume 20% polyethylene glycol (PEG)/2.5M NaCl and leaving at 4°C for 1 hour. These were spun for 15 minutes as described above and the pellets resuspended in 10 mM Tris/HCl pH 8, 1mM EDTA to 1/100th of the original volume. Residual bacteria and undissolved material were removed by spinning for 2 minutes in a microcentrifuge. Single stranded DNA for mutagenesis or sequencing was prepared from concentrated phage according to Sambrook, J., et al., 1989, supra.

Index of Examples

Example 1 Design of Insertion Point Linkers and Construction of Vectors

This example covers the construction of two derivatives of the phage vector fd-tet: a) fdTPs/Bs for the insertion of VH coding sequences; and b) fdTPs/Xh for the insertion of scFv coding sequences. The derivative vectors have a new BstEII site for insertion of sequences.

Example 2 Insertion of Immunoglobulin Fv Domain into Phage

This example covers the insertion of scFv coding sequences derived from an anti-lysozyme antibody D1.3 into fdTPs/Xh to give the construct fdTscFvD1.3.

Example 3 Insertion of Immunoglobulin VH Domain into Phage

This example covers the insertion of VH coding sequences derived from an anti-lysozyme antibody D1.3 into fdTPs/Bs to give the construct fdTVHD1.3.

Example 4 Analysis of Binding Specificity of Phage Antibodies

This example investigates the binding specificities of the constructs fdTscFvD1.3 and fdTVHD1.3.

Example 5 Construction of fdCAT2

This example covers the construction of the derivative fdCAT2 of the phage vector fdTPs/Xh. The derivative has restriction sites for enzymes that cut DNA infrequently.

Example 6 Specific Binding of Phage Antibody (pAb) to Antigen

This example shows the binding of pAb fdTscFvD1.3 to lysozyme by ELISA.

5 Example 7 Expression of FabD1.3

This example concerns the display of an antibody Fab fragment at the phage surface. The VH-CH1 chain is expressed by fdCAT2. The VL-CL chain is expressed by pUC19 in a bacterial host cell also infected with fdCAT2.

10 Example 8 Isolation of Specific, Desired Phage from a Mixture of Vector Phage

This example shows how a phage (e.g. fdTscFvD1.3) displaying a binding molecule can be isolated from vector phage by affinity techniques.

15 Example 9 Construction of pAb Expressing Anti-Hapten Activity

This example concerns the insertion of scFv coding sequences derived from the anti-oxazolone antibody NQ11 into fdTPs/Xh to generate the construct pAbNQ11. The example shows the binding of pAbNQ11 to oxazolone by ELISA.

20 Example 10 Enrichment of pAbD1.3 from Mixtures of other pAbs by Affinity Purification

This example shows how a phage (eg. pAbD1.3) displaying one sort of binding molecule can be isolated from phage (e.g. pAbNQ11) displaying another sort of binding molecule by affinity techniques.

Example 11 Insertion of a Gene Encoding an Enzyme (Alkaline Phosphate) into fdCAT2

30 This example concerns the invention of coding sequences for an enzyme into the vector fdCAT2 to give the phage enzyme, fdphoAla116.

Example 12 Measuring Enzyme Activity Phage - Enzyme

35 This example shows the functionality of an enzyme (alkaline phosphatase) when displayed at the phage surface (fdphoAla166).

Example 13 Insertion of Binding Molecules into Alternative Sites in the Phage

40 This example covers the insertion of scFv coding sequences derived from a) the anti-lysozyme antibody D1.3; and b) the anti-oxazolone antibody NQ11 into a BamH1 site of fdTPs/Xh to give the constructs fdTBam1 having an NQ11 insert.

45 Example 14 PCR Assembly of Mouse VH and VLK Repertoires for Phage Display

This example concerns a system for the display on phage of all VH and VLK repertoires encoded by a mouse. The system involves the following steps. 1) Preparation of RNA from spleen. 2) Preparation of cDNA from the RNA 3) Use of primers specific for antibody sequences to PCR amplify all VH and VLK cDNA coding sequences 4) Use of PCR to create a linker molecule from linking pairs of VH

and VLK sequences 5) Use of PCR to assemble continuous DNA molecules each comprising a VH sequence, a linker and a VLK sequence. The specific VH/VLK combination is randomly derived 6) Use of PCR to introduce restriction sites.

5

Example 15. Insertion of the Extracellular Domain of a Human Receptor for Platelet Derived Growth Factor (PDGF) Isoform BB into fdCAT2

10

This example concerns the insertion of coding sequences for the extracellular domain of the human receptor for PDGF into the vector fdCAT2 to give the construct fdhPDGFBR.

15

Example 16. Binding of 125 I-PDGF-BB to the Extracellular Domain of the Human Receptor for PDGF Isoform BB Displayed on the Surface of fd Phage. Measured using an Immunoprecipitation Assay.

20

This example shows that the human receptor PDGF Isoform BB is displayed on the surface of the phage in a form which has the ability to bind its ligand.

25

Example 17. Construction of Phagemid Containing Gene III Fused with the Coding Sequence for a Binding Molecule.

This example concerns the construction of two phagemids based on pUC119 which separately contain gene III from fdCAT2 and the gene III scFv fusion fdCAT2scFvDI.3 to generate pCAT2 and pCAT3 scFvDI.3 respectively.

30

Example 18. Rescue of Anti-Lysozyme Antibody Specificity from pCAT3scFvDI.3 by M13KO7

This example describes the rescue of the coding sequence for the gene IIIscFv fusion from pCAT3scFvDI.3 by M13MO7 helper phage growth, phage were shown to be displaying scFv anti-lysozyme activity by ELISA.

35

Example 19. Transformation Efficiency of PCAT-3 and pCAT-3 scFvDI.3 Phagemids

This example compared the efficiency of the phagemids pUC119, pCAT-3 and pCAT3scFvDI.3 and the phage fdCAT2 to transform E.coli.

40

Example 20 PCR Assembly of a Single Chain Fv Library from an Immunised Mouse

This example concerns a system for the display on phage of scFv (comprising VH and VL) from an immunised mouse using the basic technique outlined in example 14 (cDNA preparation and PCR assembly of the mouse VH and VLK repertoires) and ligating the PCR assembled sequences into fdCAT2 to create a phage library of 10^5 clones.

45

Testing of 500 clones showed that none showed specificity against phox.

Example 21. Selection of Antibodies Specific for 2-phenyl-5-oxazolone from a Repertoire from an Immunised Mouse.

50

This example shows that phage grown from the library established in example 20 can be subjected to affinity selection using phOX to select those phage displaying

scFv with the desired specificity.

Example 22. Generation of Further Antibody Specificities by the Assembly of Hierarchical Libraries.

5 This example concerns the construction of hierarchical libraries in which a given VH sequence is combined with the complete VLK repertoire and a given VLK sequence is combined with the complete VH repertoire and selection from these libraries of novel VH and VL pairings.

10 Example 23. Selection of Antibodies Displayed on Bacteriophage with Different Affinities for 2-phenyl-5-oxazolone using Affinity Chromatography

This example concerns the separation by affinity techniques of phages displaying scFv fragments with differing binding affinities for a given antigen.

15 Example 24. Construction of Phagemid pHEN1 for the Expression of Antibody Fragments Expressed on the surface of Bacteriophage following Superinfection

20 This example concerns the construction of the phagemid pHEN1 derived from pUC119. pHEN1 has the features shown in Fig. 26.

Example 25. Display of Single Chain Fv and Fab Fragments Derived from the Anti-Oxazolone Antibody NO 10.12.5 on Bacteriophage fd using pHEN1 and fdCAT2.

25 This example describes the display of scFv and Fab fragment with a specificity against phOx on the surface of a bacteriophage. For display of scFv the phagemid pHEN1 comprises the sequences encoding scFv (VH and VL) for rescue by either the phages VSM13 or fdCAT2. For display of Fab the phage fdCAT2 comprises the sequence for either the H or L chain as a fusion with g3p and the phagemid pHEN1 comprises the sequence for the appropriate H or L chain partner.

30 Example 26. Rescue of Phagemid Encoding a Gene III protein Fusion with Antibody Heavy or Light Chains by Phage Encoding the Complementary Antibody Displayed on Phage and the Use of this Technique to make Dual Combinatorial Libraries

35 This example covers the use of phage antibodies encoding the antibody heavy or light chain to rescue a phagemid encoding a gene 3 protein fusion with the complementary chain and the assay of Fab fragments displayed on phage in ELISA. The use of this technique in the preparation of a dual combinatorial library is discussed.

40 Example 27 Induction of Soluble scFv and Fab Fragments using Phagemid pHEN1

45 This example covers the generation of soluble scFv and Fab fragments from gene III fusions with sequences encoding these fragments by expression of clones in pHEN1 in an E.coli strain which does not suppress amber mutations.

50 Example 28 Increased Sensitivity in ELISA of Lysozyme

using fdTscFvD1.3 as Primary Antibody compared to Soluble scFvD1.3

5 This example covers the use of fdTscFvD1.3 in ELISA showing that lower amounts of lysozyme can be detected with phage antibody fdTscFvD1.3 than with soluble scFvD1.3.

Example 29 Direct Rescue and Expression of Mouse Monoclonal Antibodies as Single Chain Fv Fragments on the Surface of Bacteriophage fd

10 This example covers the display on phage as functional scFv fragments of two clones directly derived from cells expressing monoclonal antibodies directed against oestriol. Both clones were established to be functional using ELISA.

15 Example 30 Kinetic Properties of Alkaline Phosphatase Displayed on the Surface of Bacteriophage fd

20 This example concerns the demonstration that the kinetic properties of an enzyme, alkaline phosphatase, displayed on phage are qualitatively similar to those of the same enzyme when in solution.

Example 31 Demonstration using Ultrafiltration that Cloned Alkaline Phosphatase Behaves as Part of the Virus Particle

25 This example concerns the construction of the phage enzyme fdphoArg166 and the demonstration that both the fusion protein made and the catalytic activity observed derive from the phage particle.

Example 32 Affinity Chromatography of Phage Alkaline Phosphatase

30 This example concerns the binding of alkaline phosphatase displayed on phage to an arsenate-Sepharose affinity column and specific elution of these phage using the reaction product, phosphate.

35 Example 33 PCR Assembly of DNA Encoding the Fab Fragment of an Antibody Directed against Oxazolone

40 This example covers the construction of a DNA insert encoding a Fab fragment by separate amplification of heavy and light chain DNA sequences followed by assembly. The construct was then inserted into the phage vector fdCAT2 and the phagemid vector pHEN1 and the Fab fragment displayed on the surface was shown to be functional.

Example 34 Construction of a Gene III Deficient Helper Phage

45 This example describes the construction of a helper phage derived from M13K07 by deleting sequences in gene III. Rescue of pCAT3-scFvD1.3 is described. The scFvD1.3 is expressed at a high level as a fusion using the deletion phage, equivalent to expression using fdCAT2-scFvD1.3.

50 Example 35 Selection of bacteriophage expressing scFv fragments directed against lysozyme from mixtures according to affinity using a panning procedure

This example concerns the selection of bacteriophage according to the affinity of the scFv fragment directed against lysozyme which is expressed on their surface. The phage of different affinities were bound to Petri dishes coated with lysozyme and, following washing, bound phage eluted using triethylamine. Conditions were found where substantial enrichment could be obtained for a phage with a 5-fold higher affinity than the phage with which it was mixed.

10 Example 36 Expression of Catalytically Active Staphylococcal Nuclease on the Surface of Bacteriophage fd

This example concerns the construction of a phage enzyme which expresses Staphylococcal nuclease and the demonstration that the phage enzyme retains nuclease activity.

15 Example 37 Display of the Two Aminoterminal Domains of Human CD4 on the Surface of fd Phage

This example covers the cloning of genes for domains of CD4, a cell surface receptor and member of the immunoglobulin superfamily, into bacteriophage fd. The receptor is shown to be functional on the surface of phage by binding to the HIV protein gp120.

20 Example 38 Generation and Selection of Mutants of an Anti-4-hydroxy-3-nitrophenylacetic acid (NP) Antibody expressed on Phage using Mutator strains

This example covers the introduction of mutations into a gene for an antibody cloned in phage by growth of the phage in strains which randomly mutate DNA due to defects in DNA replication. Several mutations are introduced into phage which can then be selected from parent phage.

30 Example 39 Expression of a Fv Fragment on the Surface of Bacteriophage by Non-Covalent Association of VH and VL domains

This example shows that functional Fv fragments can be expressed on the surface of bacteriophage by non-covalent association of VH and VL domains. The VH domain is expressed as a gene III fusion and the VL domain as a soluble polypeptide. Sequences allowing expression of these domains from the anti-lysozyme antibody D1.3 in this form were introduced into phage and the resulting displayed Fv fragment shown to be functional by ELISA.

40 Example 40 A PCR Based Technique for one step Cloning of Human V-genes as Fab Constructs

This example gives methods for the assembly of Fab fragments from genes for antibodies. Examples are given for genes for antibodies directed against Rhesus-D in a human hybridoma and a polyclonal lymphoblastic cell line.

50 Example 41 Selection of Phage Displaying a Human Fab Fragment directed against the Rhesus-D Antigen by binding to Cells displaying the Rhesus D Antigen on their Surface

This example concerns the construction of, and display of phage antibodies from, a phagemid encoding a human Fab fragment directed against the Rhesus D antigen. Phage displaying this antigen were then affinity selected from a background of phage displaying scFvD1.3 anti-lysozyme on the basis of binding to Rhesus-D positive red blood cells.

Example 42 A PCR Based Technique for One Step Cloning of Human scFv Constructs

This example describes the generation of libraries of scFv fragments derived from an unimmunized human. Examples are given of the preparation for phage display of libraries in phagemids of scFv fragments derived from IgG and IgM sequences.

Example 43 Isolation of Binding Activities from a Library of scFvs from an Unimmunized Human

This example describes the isolation, from the library of scFv fragments derived from IgM genes of an unimmunized human, of clones for phage antibodies directed against BSA, lysozyme and oxazolone. Selection was by panning or affinity chromatography and analysis of binding specificity by ELISA. Sequencing of the clones showed them to be of human origin.

Example 44 Rescue of human IgM library using helper phage lacking gene 3 (g3)

This example covers the isolation, from the library of scFv fragments of unimmunized human IgM genes, of clones of phage antibodies of clones for phage antibodies specific for thyroglobulin and oxazolone. In this example rescue was with M13K07gIII No3 (NCTC12478), a helper phage defective in gene III. Fewer rounds of selection appeared necessary for a phagemid library rescued with this phage compared to one rescued with M13K07.

Example 45 Alteration of Fine Specificity of scFvD1.3 displayed on Phage by Mutagenesis and Selection on Immobilized Turkey Lysozyme

This example covers the in vitro mutagenesis of pCATscFvD1.3 by replacement, with random amino acids, of residues known to be of importance in the preferential recognition of hen egg lysozyme over turkey egg lysozyme by scFvD1.3. Following selection for phage antibodies recognising turkey egg lysozyme by affinity chromatography, clones were analysed for specificity by ELISA. Two groups of clones were found with more equal recognition of hen and turkey lysozymes, one with increased ELISA signal with the turkey enzyme and one with reduced signal for the hen enzyme.

Example 46 Modification of the Specificity of an Antibody by Replacement of the VLK Domain by a VLK Library derived from an Unimmunised Mouse

This example shows that replacement of the VL domain

of scFvD1.3 specific for hen eggwhite lysozyme (HEL) with a library of VL domains allows selection of scFv fragments which bind also to turkey eggwhite lysozyme (TEL). The scFv fragments were displayed on phage and selection by panning on tubes coated with TEL. Analysis by ELISA showed clones with enhanced binding to TEL compared to HEL. Those with highest binding to TEL were sequenced.

10 Example 47 Selection of a Phage Antibody Specificity by binding to an Antigen attached to Magnetic Beads. Use of a Cleavable Reagent to allow Elution of Bound Phage under Mild Conditions

15 This examples covers the use of a cleavable bond in the affinity selection method to allow release of bound phage under mild conditions. pAbN011 was enriched approximately 600 fold from a mixture with pAbD1.3 by selection using biotinylated Ox-BSA bound to magnetic beads. The cleavage of a bond between BSA and the biotin allows elution of the phage.

20 Example 48 Use of Cell Selection to provide an Enriched Pool of Antigen Specific Antibody Genes, Application to reducing the Complexity of Repertoires of Antibody Fragments Displayed on the Surface of Bacteriophage

25 This example covers the use of cell selection to produce an enriched pool of genes encoding antibodies directed against 4-hydroxy-3-nitrophenylacetic acid and describes how this technique could be used to reduce the complexity of antibody repertoires displayed on the surface of bacteriophage.

30 Example 1

Design of Insertion Point Linkers and Construction of Vectors

35 The vector fd-tet has two BstEII restriction sites flanking the tetracycline resistance gene (fig 3). Since the strategy for inserting the VH fragments was to ligate them into a newly inserted BstEII site within gene III, it was advantageous to delete the original BstEII sites from fd-tet. This was achieved by digesting fd-tet with the restriction enzyme BstEII, filling-in the 5' overhangs and re-ligating to generate the vector fdT6Bst. Digestion of fd-tet with BstEII (0.5 units/ μ l) was carried out in 1x KGB buffer (100 mM potassium glutamate, 23 mM Tris-acetate (pH 7.5), 10 mM magnesium acetate, 50 μ g/ml bovine serum albumin, 0.5 mM dithiothreitol (Sambrook, J., et al., 1989, supra.) with DNA at a concentration of 25 ng/ μ l. The 5' overhang was filled in, using 2x KGB buffer, 250 μ M each dNTP's (Pharmacia Ltd., Pharmacia House, Midsummer Boulevard, Milton Keynes, Bucks., UK.) and Klenow Fragment (Amersham International, Lincoln Place, Green End, Aylesbury, Bucks., UK) at 0.04 units/ μ l. After incubating for 1 hour at room temperature, DNA was extracted with

phenol/chloroform and precipitated with ethanol.

Ligations were carried out at a DNA concentration of 50ng/ μ l). Ligations were transformed into competent TG1 cells and plated onto TY plates supplemented with 15 μ g/ml tetracycline. This selects for vectors where the gene for tetracycline resistance protein has reinserted into the vector during the ligation step. Colonies were picked into 25 mls of 2xTY medium supplemented with 15 μ g/ml tetracycline and grown overnight at 37°C.

Double stranded DNA was purified from the resulting clones using the gene-clean II kit (Bio101 Inc., PO Box 2284, La Jolla, California, 92038-2284, USA.) and according to the small scale rapid plasmid DNA isolation procedure described therein. The orientation of 5 of the resulting clones was checked using the restriction enzyme ClaI. A clone was chosen which gave the same pattern of restriction by ClaI as fd-tet, but which had no BstE II sites.

In vitro mutagenesis of fdT δ Bst was used to generate vectors having appropriate restriction sites that facilitate cloning of antibody fragments downstream of the gene III signal peptide and in frame with the gene III coding sequence. The oligonucleotide directed mutagenesis system version 2 (Amersham International) was used with oligo 1 (figure 4) to create fdTPs/Bs (to facilitate cloning of VH fragments). The sequence of fdTPs/Bs (figure 4) was confirmed using the sequenase version 2.0 kit (USB Corp., PO Box 22400, Cleveland, Ohio, 44122, USA.) with oligo 3 (figure 4) as a primer.

A second vector fdTPs/Xh (to facilitate cloning of single chain Fv fragments) was generated by mutagenising fdTPs/Bs with oligo 2 according to the method of Venkitaraman, A.R., Nucl. Acid Res. 17, p 3314. The sequence of fdTPs/Xh (figure 4) was confirmed using the sequenase version 2.0 kit (USB Corp.) with oligo 3 as a primer.

Clearly, alternative constructions will be apparent to those skilled in the art. For example, M13 and/or its host bacteria could be modified such that its gene III could be disrupted without the onset of excessive cell death; the modified fd gene III, or other modified protein, could be incorporated into a plasmid containing a single stranded phage replication origin, such as pUC119, superinfection with modified phage such as K07 would then result in the encapsulation of the phage antibody genome in a coat partially derived from the helper phage and partly from the phage antibody gene III construct.

The detailed construction of a vector such as fdTPs/Bs is only one way of achieving the end of a phage antibody. For example, techniques such as sticky feet cloning/mutagenesis (Clackson, T. and Winter, G. 1989

Nucl. Acids. Res., 17, p 10163-10170) could be used to avoid use of restriction enzyme digests and/or ligation steps.

Example 2.

5 Insertion of Immunoglobulin Fv Domain into Phage

10 The plasmid scFv D1.3 myc (gift from G. Winter and A. Griffiths) contains VH and VL sequences from the antibody D1.3 fused via a peptide linker sequence to form a single chain Fv version of antibody D1.3. The sequence of the scFv and surrounding sequences in scFvD1.3 myc is shown in figure 5.

15 The D1.3 antibody is directed against hen egg lysozyme (Harper, M. et al., 1987, Molec. Immunol. 24, 97-108) and the scFv form expressed in E.coli has the same specificity (A. Griffiths and G. Winter personal Communication).

20 Digestion of scFv D1.3 myc with PstI and XhoI (these restriction sites are shown on Fig. 5), excises a fragment of 693 bp which encodes the bulk of the scFv. Ligation of this fragment into fdTPs/Xh cleaved with PstI and XhoI gave rise to the construct fdTscFvD1.3 encoding the gene III signal peptide and first amino acid fused to the complete D1.3 scFv, followed by the mature gene III protein from amino acid 2.

25 The vector fdTPs/Xh was prepared for ligation by digesting with the PstI and XhoI for 2 hours followed by digestion with calf intestinal alkaline phosphatase (Boehringer Mannheim UK Ltd., Bell Lane, Lewes, East Sussex, BN7 1LG) at one unit/ul for 30 minutes at 37°C. Fresh calf intestinal alkaline phosphatase was added to a final total concentration of 2 units/ul and incubated for a further 30 minutes at 37°C. The reaction was extracted three times with phenol/chloroform, precipitated with ethanol and dissolved in water. The insert from scFvD1.3 myc was excised with the appropriate restriction enzymes (PstI and XhoI) extracted twice with phenol/chloroform, precipitated with ethanol and dissolved in water. Ligations were carried out as described in example 1, except both vector and insert samples were at a final concentration of 5 ng/ul each. The formation of the correct construct was confirmed by sequencing as described in example 1.

45 To demonstrate that proteins of the expected size were produced, virions were concentrated by PEG precipitation as described above. The samples were prepared for electrophoresis as described in Sambrook J. et al 1989 supra. The equivalent of 2mls of supernatant was loaded onto an 18% SDS polyacrylamide gel. After electrophoresis, the gel was soaked in gel running buffer (50 mM Tris, 380 mM Glycine, 0.1% SDS) with 20% methanol for 15 minutes. Transfer to nitrocellulose filter was executed in fresh 1x running buffer/20% methanol using

TE70 Semi Phor a semi-dry blotting apparatus (Hoeffer, 654 Minnesota Street, Box 77387, San Francisco, California 94107, USA.).

After transfer, the filter was blocked by incubation for 1 hour in a 2% solution of milk powder (Marvel) in phosphate buffered saline (PBS). Detection of scFv and VH protein sequences in the phage antibody fusion proteins was effected by soaking the filter for 1 hour with a 1/1000 dilution (in 2% milk powder) of a rabbit polyclonal antiserum raised against affinity purified, bacterially expressed scFv fragment (gift from G. Winter). After washing with PBS (3 x 5 minute washes), bound primary antibody was detected using an anti-rabbit antibody conjugated to horseradish peroxidase (Sigma, Fancy Road, Poole, Dorset, BH17 7NH, UK.) for 1 hour. The filter was washed in PBS/0.1% triton X-100 and developed with 0.5 mg/ml 3,3'-diaminobenzidine tetrahydrochloride (DAB), 0.02% cobalt chloride, 0.03% hydrogen peroxide in PBS.

The results showed that with clones fdTVHD1.3 (from example 3 incorporating sequences coding for VH) and fdTscFvD1.3 (incorporating sequences coding for scFv) a protein of between 69,000 and 92,500 daltons is detected by the anti-Fv serum. This is the expected size for the fusion proteins constructed. This product is not observed in supernatants derived from fd-tet, fdT6Bst or fdTPs/Kh.

Example 3.

Insertion of Immunoglobulin VH Domain into Phage Antibody

The VH fragment from D1.3 was generated from the plasmid pSW1-VHD1.3-TAG1 (Ward, E.S. et al., 1989 supra.). Digestion of this plasmid with PstI and BstEII generates the fragment shown between positions 113 and 432 in figure 5. Cloning of this fragment into the PstI and BstEII sites of fdTPs/Bs gave rise to the construct fdTVHD1.3 which encodes a fusion protein with a complete VH domain inserted between the first and third amino acids of the mature gene III protein (amino acid two has been deleted).

The methods used were exactly as in example 2 except that the vector used was fdTPs/Bs digested with PstI and BstEII.

Example 4.

Analysis of Binding Specificity of Phage Antibodies

The binding of the various phage antibodies to the specific antigen, lysozyme, was analysed using ELISA techniques. Phage antibodies (e.g. fdTVHD1.3 and fdTsc/FvD1.3) were grown in E.coli and Phage antibody particles were precipitated with PEG as described in the materials and methods. Bound phage antibody particles were detected using polyclonal sheep serum raised against the closely related phage M13.

ELISA plates were prepared by coating 96 well plates (Falcon Microtest III flexible plate. Falcon: Becton Dickinson Labware, 1950 Williams Drive, Oxnard, California, 93030, USA.) with 200 μ l of a solution of lysozyme (1mg/ml unless otherwise stated) in 50 mM NaHCO₃ for 16-24 hours. Before use, this solution was removed, the plate rinsed several times in PBS and incubated with 200 μ l of 2% milk powder/PBS for 1 hour. After rinsing several times with PBS, 100 μ l of the test samples were added and incubated for 1 hour. Plates were washed (3 rinses in 0.05% Tween 20/PBS followed by 3 rinses in PBS alone). Bound phage antibodies were detected by adding 200 μ l/well of a 1/1000 dilution of sheep anti-M13 polyclonal antiserum (gift from G. Winter, although an equivalent antibody can be readily made by one skilled in the art using standard methodologies) in 2% milk powder/PBS and incubating for 1 hour. After washing as above, plates were incubated with biotinylated anti-sheep antibody (Amersham International) for 30 minutes. Plates were washed as above, and incubated with streptavidin-horseradish peroxidase complex (Amersham International). After a final wash as above, 0.5 mg/ml ABTS substrate in citrate buffer was added (ABTS = 2,2'-azino-bis (3-ethylbenzthiazoline sulphonic acid); citrate buffer = 50 mM citric acid, 50 mM tri-sodium citrate at a ratio of 54:46. Hydrogen peroxide was added to a final concentration of 0.003% and the plates incubated for 1 hour. The optical density at 405 nm was read in a Titertek multiskan plate reader.

Figure 6 shows the effect of varying the amount of phage antibody. 100 μ l of various dilutions of PEG precipitated phage were applied and the amount expressed in terms of the original culture volume from which it was derived. Signals derived from both the scFv containing phage antibody (fdTscFvD1.3) and the VH containing phage antibody (fdTVHD1.3) and the VH containing phage antibody were higher than that derived from the phage antibody vector (fdTPs/Xh). The highest signal to noise ratio occurs using the equivalent of 1.3 mls of culture.

Figure 7 shows the results of coating the plates with varying concentrations of lysozyme or bovine serum albumin (BSA). The equivalent of 1 ml of the original phage antibody culture supernatant was used. The signals from supernatants derived from fdTscFvD1.3 were again higher than those derived from fdTPs/Xh when lysozyme coated wells were used. There was no significant difference between these two types of supernatant when the plates were coated with BSA. Broadly speaking the level of signal on the plates is proportional to the amount of lysozyme coated. These results demonstrate that the binding detected is specific for lysozyme as the antigen.

Example 5:Construction of fd CAT 2

It would be useful to design vectors that enable the use of restriction enzymes that cut DNA infrequently, thus avoiding unwanted digestion of the antibody gene inserts within their coding sequence. Enzymes with an eight base recognition sequence are particularly useful in this respect, for example NotI and SfiI. Chaudhary et al (PNAS 87 p1066-1070, 1990) have identified a number of restriction sites which occur rarely in antibody variable genes. The applicant has designed and constructed a vector that utilises two of these sites, as an example of how this type of enzyme can be used. Essentially sites for the enzymes ApaI and NotI were engineered into fdTPS/Xh to create fdCAT2.

The oligonucleotide:

5'ACT TTC AAC AGT TTC TGC GGC CGC CCG TTT GAT CTC GAG CTC CTG CAG TTG GAC CTG TGC ACT GTG AGA ATA GAA 3'

was synthesised (supra fig 4 legend) and used to mutagenise fdTPS/Xh using an in vitro mutagenesis kit from Amersham International as described in example 1, to create fd-CAT2. The sequence of fd-CAT2 was checked around the site of manipulation by DNA sequencing. The final sequence around the insertion point within gene III is shown in figure 8.

N.B. fdCAT2 is also referred to herein by the alternative terminologies fd-tet-DOG1 and fdDOG1.

Example 6Specific Binding of Phage-antibody (pAb) to Antigen

The binding of pAb D1.3 (fdTscFvD1.3 of example 2) to lysozyme was further analysed by ELISA.

Methods.1. Phage growth.

Cultures of phage transduced bacteria were prepared in 10-100 mls 2 x TY medium with 15 µg/ml tetracycline and grown with shaking at 37°C for 16-24 hrs. Phage supernatant was prepared by centrifugation of the culture (10 min at 10,000 rpm, 8 x 50 ml rotor, Sorval RC-5B centrifuge). At this stage, the phage titre was 1 - 5 x 10¹⁰/ml transducing units. The phage were precipitated by adding 1/5 volume 20% PEG 2.5 M NaCl, leaving for 1 hr at 4°C, and centrifuging (supra). The phage pellets were resuspended in 10 mM Tris-HCl, 1mM EDTA pH 8.0 to 1/100th of the original volume, and residual bacteria and aggregated phage removed by centrifugation for 2 min in a bench microcentrifuge.

ELISA

Plates were coated with antigen (1 mg/ml antigen) and blocked as described in example 4. 2 x 10¹⁰ phage transducing units were added to the antigen coated plates in phosphate buffered saline (PBS) containing 2% skimmed milk powder (MPBS). Plates were washed between each step

with three rinses of 0.5% Tween-20 in PBS followed by three rinses of PBS. Bound phage was developed by incubating with sheep anti-M13 antisera and detected with horseradish peroxidase (HRP) conjugated anti-goat serum (Sigma, Poole, Dorset, UK) which also detects sheep immunoglobulins and ABTS (2,2'-azino-bis (3-ethylbenzthiazoline sulphonic acid). Readings were taken at 405 nm after a suitable period. The results (figure 9) show that the antibody bearing-phage had the same pattern of reactivity as the original D1.3 antibody (Harper, M., Lema, F., Boulot, G., and Poljak, F.J. (1987) *Molec. Immunol.* 24, 97-108), and bound to hen egg-white lysozyme, but not to turkey egg-white lysozyme, human lysozyme or bovine serum albumin. The specificity of the phage is particularly illustrated by the lack of binding to the turkey egg-white lysozyme that differs from hen egg-white lysozyme by only 7 amino acids.

Example 7.

Expression of Fab D1.3

The aim of this example was to demonstrate that the scFv format used in example 2 was only one way of displaying antibody fragments in the pAb system. A more commonly used antibody fragment is the Fab fragment (figure 1) and this example describes the construction of a pAb that expresses a Fab-like fragment on its surface and shows that it binds specifically to its antigen. The applicant chose to express the heavy chain of the antibody fragment consisting of the VH and CH1 domains from coding sequences within the pAb itself and to co-express the light chain in the bacterial host cell infected with the pAb. The VH and CH1 regions of anti-lysozyme antibody D1.3 were cloned in fd CAT2, and the corresponding light chain cloned in plasmid pUC19. The work of Skerra and Pluckthun (Science 240, p1038-1040 (1988) and Better et al 1988 *supra*; demonstrated that multimeric antigen binding fragments of the antibody molecule could be secreted into the periplasm of the bacterial cell in a functional form using suitable signal sequences. However, in these publications, special measures were described as being needed to recover the binding protein from the cell, for example Skerra and Pluckham needed to recover the Fv fragment from the periplasm by affinity chromatography. The present applicants have shown that it is possible to direct the binding molecule to the outside of the cell on a phage particle, a process that requires several events to occur: correct secretion and folding of the binding molecule; association of the chains of the binding molecule; correct assembly of the phage particle; and export of the intact phage particle from the cell.

Alternatively, it is possible however, to express the light chain from within the pAb genome by, for

example, cloning an expression cassette into a suitable place in the phage genome. Such a suitable place would be the intergenic region which houses the multicloning sites engineered into derivative of the related phage M13 (see, for example, Yanisch-Perron, C. et al., Gene 33, p103-119, (1985)).

The starting point for this example was the clone Fab D1.3 in pUC19, a map of which is shown in figure 10. The regions hybridising with the oligonucleotides KSJ6 and 7 below are shown underlined in fig 10. The sequence encoding the VH-CH1 region (defined at the 5' and 3' edges by the oligonucleotides KSJ6 and 7 below) was PCR amplified from Fab D1.3 in pUC19 using oligonucleotides KSJ 6 and 7, which retain the Pst I site at the 5' end and introduce a Xho I site at the 3' end, to facilitate cloning into fd CAT2. The sequences for the oligonucleotides KSJ6 and 7 are shown below. The underlined region of KSJ7 shows the portion hybridising with the sequence for D1.3.

KSJ6: 5' AGG TGC AGC TGC AGG AGT CAG G 3'

KSJ7: 5' GGT GAC CTC GAG TGA AGA TTT GGG CTC AAC TTT C 3'

PCR conditions were as described in example II, except that thirty cycles of PCR amplification were performed with denaturation at 92°C for 45 seconds, annealing at 55°C for 1 minute and extension at 72°C for 1 minute. The template used was DNA from TG1 cells containing Fab D1.3 in pUC19 resuspended in water and boiled. The template DNA was prepared from the colonies by picking some colony material into 100µl of distilled H₂O and boiling for 10 mins. 1µl of this mixture was used in a 20µl PCR. This regime resulted in amplification of the expected fragment of approximately 600bp. This fragment was cut with Pst I and Xho I, purified from an agarose gel and ligated into Pst I/Xho I-cut fdCAT2. The PCR mixture was extracted with phenol/chloroform and ethanol precipitated (Sambrook et al. supra.) before digestion with PstI and XhoI (New England Biolabs according to manufacturers recommendations. The fragment was resolved on 1% Tris-Acetate EDTA agarose gel (Sambrook et al. supra) and purified using Geneclean (BIO 101, Geneclean, La Jolla, San Diego, California, USA) according to manufacturers recommendations.

fd-CAT2 vector DNA was digested with Pst I and Xho I (New England BioLabs) according to manufacturers recommendations, extracted with phenol/chloroform and ethanol precipitated (Sambrook et al. supra.).

75ng of Pst I/Xho I-digested vector DNA was ligated to 40ng of PCR-amplified PstI /Xho I-digested hEGF-R fragment in 12µl of ligation buffer (66mM TrisHCl (pH7.6), 5mM MgCl₂, 5mM dithiothreitol, (100µg/ml bovine serum albumin, 0.5mM ATP, 0.5mM Spermidine) and 40C units T4 DNA ligase (New England BioLabs) for 16 hours at 16°C.

Two μ l of the ligation mixture was transformed into 200 μ l of competent E.coli MC1061 cells, plated on 2TY agar containing 15 μ g/ml tetracycline and incubated at 30°C for 20 hours. A portion of the ligation reaction mixture was transformed into E.coli MC1061 (Available from, for example Clontech Laboratories Inc, Palo Alto, California) and colonies identified by hybridisation with the oligonucleotide D1.3CDR3A as described in example 10. The presence of the VHCH1 gene fragment was likewise confirmed by PCR, using oligonucleotides KSJ6 and 7. A representative clone was called fd CAT2VHCH1 D1.3. The heavy chain was deleted from Fab D1.3 in pUC19 by Sph I cleavage of Fab D1.3 plasmid DNA. The pUC 19 2.7Kb fragment containing the light chain gene was purified from a TAE agarose gel, and 10ng of this DNA self-ligated and transformed into competent E.coli TG1. Cells were plated on 2TY agar containing ampicillin (100 μ g/ml) and incubated at 30°C overnight. The resulting colonies were used to make miniprep DNA (Sambrook et al. supra), and the absence of the heavy chain gene confirmed by digestion with Sph I and Hind III. A representative clone was called LCD1.3 DHC.

An overnight culture of fd CAT2VHCH1 D1.3 cells was microcentrifuged at 13,000Xg for 10 minutes and 50 μ l of the supernatant containing phage particles added to 50 μ l of an overnight culture of LCD1.3 DHC cells. The cells were incubated at 37°C for 10 minutes and plated on 2TY agar containing ampicillin (100 μ g/ml) and 15 μ g/ml tetracycline. Phage were prepared from some of the resulting colonies and assayed for their ability to bind lysozyme as described in example 6.

The results (Figure 11) showed that when the heavy and light chain Fab derivatives from the original antibody D1.3 were present, the pAb bound to lysozyme. pAb expressing the fd VHCH1 fragment did not bind to lysozyme unless grown in cells also expressing the light chain. This shows that a functional Fab fragment was produced by an association of the free light chain with VHCH1 fragment fused to gene III and expressed on the surface of the pAb.

Example 8

Isolation of Specific, Desired Phage from a Mixture of Vector Phage.

The applicant purified pAb (D1.3) (originally called fdTscFvD1.3 in example 2) from mixtures using antigen affinity columns. pAb (D1.3) was mixed with vector fd phage (see table 1) and approximately 10^{12} phage passed over a column of lysozyme-Sepharose (prepared from cyanogen bromide activated sepharose 4B (Pharmacia, Milton Keynes, Bucks, UK.) according to the manufacturers instructions. TG1 cells were infected with appropriate dilutions of the elutes and the colonies derived, were

analysed by probing with an oligonucleotide that detects only the pAb (D1.3) see Table 1 and Fig. 12. A thousand fold enrichment of pAb(D1.3) was seen with a single column pass. By growing the enriched phage and passing it down the column again, enrichments of up to a million fold were seen.

Enrichment was also demonstrated using purely immunological criteria. For example, 10^{12} phage (at a ratio of 1 pAb (D1.3) to 4×10^6 fdTPs/Bs) was subjected to two rounds of affinity selection, and then 26 colonies picked and grown overnight. The phage was then assayed for lysozyme binding by ELISA (as example 6). Five colonies yielded phage with lysozyme binding activities, see table 1, and these were shown to encode the scFv (D1.3) by PCR screening (example 13, using 30 cycles of 1 minute at 92°C , 1 minute at 60°C , 1 minute at 72°C using CDR3PCR1 and oligo 3 (fig. 4) as primers).

Thus very rare pAbs can be fished out of large populations, by using antigen to select and then screen the phage.

In this example, affinity chromatography of pAbs and oligonucleotide probing were carried out as described below.

Approximately 10^{12} phage particles in 1ml MPBS were loaded onto a 1 ml lysozyme-Sepharose affinity column which had been prewashed in MPBS. The column was washed in turn with 10 ml PBS; then 10 ml 50 mM Tris-HCl, 500 mM NaCl pH 7.5; then 10ml 50 mM Tris-HCl 500 mM NaCl pH 8.5; then 5 mls 50 mM Tris-HCl, 500 mM NaCl pH 9.5 (adjusted with triethylamine) and then eluted with 5 ml 100 mM triethylamine. The eluate was neutralised with 0.5 M sodium phosphate buffer pH 6.8 and the phage plated for analysis. For a second round of affinity chromatography, the first column eluate was plated to about 30,000 colonies per petri dish. After overnight growth, colonies were then scraped into 5 ml $2 \times$ TY medium, and a 20 μl aliquot diluted into 10 ml fresh medium and grown overnight. The phage was PEG precipitated as described above, resuspended in 1 ml MPBS and loaded onto the column, washed and eluted as above.

Oligonucleotides synthesised:

CDR3PCR1 5'TGA GGA C(A or T) C(A or T) GC CGT CTA CTA CTG TGC 3'

40 pmole of oligonucleotide VH1FOR (Ward, E. S., et al (1989) Nature 341, 544-546), specific to pAb (D1.3) was phosphorylated with 100 μCi α - ^{32}P ATP, hybridised (1pmole/ml) to nitrocellulose filters at 67°C in $6 \times$ saline sodium citrate (SSC) Sambrook et al., supra. buffer for 30 minutes and allowed to cool to room temperature for 30 mins, washed 3×1 min at 60°C in $0.1 \times$ SSC.

Example 9

Construction of pAb Expressing Anti-hapten Activity

Oxazolone is a hapten that is commonly used for studying the details of the immune response. The anti-oxazolone antibody, NQ11 has been described previously (E. Gherardi, R. Pannell, C. Milstein, J. Immunol. Method 126 61-68). A plasmid containing the VH and VL gene of NQ11 was converted to a scFv form by inserting the BstEII/SacI fragment of scFvD1.3 myc (nucleotides 432-499 of Fig. 5) between the VH and VL genes to generate pscFvNQ11, the sequence of which is shown in fig. 13. This scFv was cloned into the PstI/XhoI site of FdTPs/Xh (as described earlier) to generate pAb NQ11 has an internal PstI site and so it was necessary to do a complete digest of pscFvNQ11 with XhoI followed by a partial digest with PstI).

The specific binding of pAb NQ11 was confirmed using ELISA. ELISA plates were coated at 37°C in 50 mM NaHCO₃ at a protein concentration of 200 µg/ml. Plates were coated with either hen egg lysozyme (HEL), bovine serum albumin (BSA), or BSA conjugated to oxazolone (OX-BSA) (method of conjugation in Makela O., Kartinen M., Pelkonen J.L.T., Karjalainen K. (1978) J. Exp. Med. 148 1644). Preparation of phage, binding to ELISA plates, washing and detection was as described in example 6. Samples were assayed in duplicate and the average absorbance after 10 minutes presented in figure 14.

This result demonstrates that the pAb NQ11 binds the correct antigen. Figure 14 also shows that pAb D1.3 and pAb NQ11 bind only to the antigen against which the original antibodies were raised.

Example 10

Enrichment of pAb D1.3 from Mixtures of Other pAb by Affinity Purification

3×10^{10} phage in 10 mls of PBSM at the ratios of pAb D1.3 to pAb NQ11 shown in table 2 were passed over a 1 ml lysozyme Sepharose column. Washing, elution and other methods were as described in example 8 unless otherwise stated. Eluates from the columns were used to infect TG1 cells which were then plated out. Colonies were probed with a probe which distinguishes pAb D1.3 from pAb NQ11. The sequence of this oligonucleotide (D1.3CDR3A) is:-

5'GTA GTC AAG CCT ATA ATC TCT CTC 3'

Table 2 presents the data from this experiment. An enrichment of almost 1000 fold was achieved in one round and an enrichment of over a million fold in two rounds of purification. This parallels the result described in example 8.

Example 11

Insertion of a Gene Encoding an Enzyme (Alkaline phosphatase) into fd-CAT2

As an example of the expression of a functional

enzyme on the bacteriophage surface, the applicants have chosen bacterial alkaline phosphatase, an enzyme that normally functions as a dimer (McCracken, S. and Meighen, E., J. Biol. Chem. 255, p2396-2404, (1980)). The oligonucleotides were designed to generate a PCR product with an *Apa* I site at the 5' end of *phoA* gene and a *Not* I site at its 3' end, thus facilitating cloning into *fd*-CAT 2 to create a gene III fusion protein. The oligonucleotides synthesised were:

10 *phoA*1:5' TAT TCT CAC AGT GCA CAA ACT GTT GAA CGG ACA CCA
GAA ATG CCT GTT CTG 3' and,
*phoA*2:5' ACA TGT ACA TGC GGC CGC TTT CAG CCC CAG AGC GGC
TTT C3'

15 The sequence of the *phoA* gene is presented in Chang C. N. et al., Gene 44, p121-125 (1986). The plasmid amplified (*pEK86*) contains an alkaline phosphate gene which differs from the sequence of Chang et al, by a mutation which converts arginine to alanine at position 166.

20 The PCR reaction was carried out in 100µl of 10 mM Tris/HCl pH 8.3, containing 50 mM KCl, 5mM dNTP 2.5 mM MgCl₂, 0.01% gelatin, 0.25 units/µl of Taq polymerase (Cetus/Perkin Elmer) and 0.5µg/ml template. The template was the *pEK86* plasmid (described by Chaidaroglou et al., Biochemistry 27 p8338-8343, 1988). The PCR was carried
25 out in a Techne (Techne, Duxford, Cambridge, UK) PHC-2 dri-block using thirty cycles of 1 min at 92°C, 2 min at 50°C, 3 min at 72°C.

The resultant product was extracted with phenol:chloroform, precipitated with ethanol, and the
30 pellet dissolved in 35µl water. Digestion with 0.3 units/µl of *Apa* I was carried out in 150µl volume according to manufacturers instructions for two hours at 37°C. After heat inactivation of the enzyme at 65°C, NaCl was added to a final concentration of 150mM and 0.4
35 units/µl *Not*I enzyme added. After incubation for 2 hours at 37°C, the digest was extracted with phenol:chloroform and precipitated as above, before being dissolved in 30µl of water. The vector *fd*-CAT2 was sequentially digested with *Apa* I and *Not*I according to the manufacturers
40 instructions and treated with calf intestinal alkaline phosphatase as described in example 2. The sample was extracted three times with phenol:chloroform, precipitated with ethanol and dissolved in water. The ligations were performed with a final DNA concentration
45 of 1-2ng/µl of both the cut *fd*-CAT2 and the digested PCR product. The ligations were transformed into competent TGI cells and plated on 2xTY tet plates. Identification of clones containing the desired insert was by analytical PCR performed using the conditions and primers above, on
50 boiled samples of the resulting colonies. The correct clone containing the *phoA* gene fused in frame to gene III was called *fd-phoA* 166. The sequence at the junction

of the cloning region is given in figure 15.

Example 12

Measuring Enzyme Activity of Phage-enzyme

5 Overnight cultures of TGI or KS272 (E.coli cells lacking phoA. Strauch K. L., and Beckwith J. PNAS 85 1576-1580, 1988) cells containing either fd-phoA1a 166 or fd-CAT2 were grown at 37°C in 2xTY with 15µg/ml tetracycline. Concentrated, PEG precipitated phage were prepared as described earlier. Enzyme assays (Malamy, 10 M.H. and Horecker B.L., Biochemistry 3, p1893-1897, (1964)) were carried out at 24°C in a final concentration of 1M Tris/HCl pH 8.0, 1mM 4-nitrophenyl phosphate (Sigma), 1mM MgCl₂. 100µl of a two times concentrate of this reaction mixture was mixed with 100µl of the test 15 sample in a 96 well plate. Absorbance readings were taken every minute for 30 minutes at a wavelength of 405nm in a Titretek Mk 2 plate reader. Initial reaction rates were calculated from the rate of change of absorbance using a molar absorbance of 17000 l/mol/cm.

20 Standard curves (amount of enzyme vs. rate of change of absorbance) were prepared using dilutions of purified bacterial alkaline phosphatase (Sigma type III) in 10mM Tris/HCl pH 8.0, 1mM EDTA. The number of enzyme molecules in the phage samples were estimated from the 25 actual rates of change of absorbance of the phage samples and comparison to this standard curve.

The results in Table 3 show that alkaline phosphatase activity was detected in PEG precipitated material in the sample containing fd-phoA1a166 but not 30 fd-CAT2. Furthermore, the level of activity was consistent with the expected number of 1-2 dimer molecules of enzyme per phage. The level of enzyme activity detected was not dependent on the host used for growth. In particular, fd-phoA1a166 grown on phoA minus 35 hosts showed alkaline phosphatase activity.

Therefore, the phage expressed active alkaline phosphatase enzyme, from the phoA-gene III fusion, on the phage surface.

Example 13

Insertion of Binding Molecules into Alternative Sites in the Phage

The availability of an alternative site in the phage for the insertion of binding molecules would open up the possibility of more easily expressing more than one 45 binding molecule e.g. an antibody fragment in a single pAb. This may be used to generate single or multiple binding specificities. The presence of two distinct binding activities on a single molecule will greatly increase the utility and specificity of this molecule. 50 It may be useful in the binding of viruses with a high mutational rate such as human immunodeficiency virus. In addition, it may be used to bring antigens into close

proximity (e.g. drug targetting or cell fusion) or it may act as a "molecular clamp" in chemical, immunological or enzymatic processes.

5 The vector fd-tet and the derivatives described here, have a single BamHI site in gene 3. This has previously been used for the expression of peptide fragments on the surface of filamentous bacteriophage (Smith GP. (1985) Science 228 p1315-1317 and de la Cruz et al. (1988) J Biol. Chem. 263 p4318-4322). This provides a potential alternative site for the insertion of antibody fragments.

DNA fragments encoding scFv's from D1.3 or NQ11 were generated by PCR using the primers shown below. These primers were designed to generate a fragment with BamHI sites near both the termini, to enable cloning into the BamHI site of gene3 (see figure 16(1)). The oligonucleotides used, also ensure that the resulting PCR product lacks PstI and XhoI restriction sites normally used for manipulating the scFv's (see figure 16(1)). This will facilitate subsequent manipulation of a second antibody fragment in the usual way at the N terminus of gene 3. The oligonucleotides used were:-
 G3Bam1 5' TTT AAT GAG GAT CCA CAG GTG CAG CTG CAA GAG 3'
 G3Bam2 5' AAC GAA TGG ATC CCG TTT GAT CTC AAG CTT 3'.

25 Preparation of vector and PCR insert

The PCR reaction was carried out in an 80 µl reaction as described in example 11 using 1ng/µl of template and 0.25U/µl of Taq polymerase and a cycle regime of 94°C for 1 minute, 60°C for 1 minute and 70°C for 2 minutes over 30 cycles. The template was either pscFvNQ11 (example 9) or scFvD1.3 myc (example 2). Reaction products were extracted with phenol:chloroform, precipitated, dissolved in water and digested with BamHI according to manufacturers instructions. The digest was re-extracted with phenol: chloroform, precipitated and dissolved in water.

The vector fdTPs/Xh was cleaved with BamHI and treated with calf intestinal phosphatase and purified as described in example 2. Ligations were set up at a vector concentration of approximately 6ng/µl and a PCR insert concentration of approximately 3ng/µl. These were ligated for 2.5 hours at room temperature before transforming into competent TGI cells and plating on TY tet plates. The resultant colonies were probed as described in example 8. DNA was prepared from a number of colonies and the correct orientation and insert size confirmed by restriction digestion with Hind III in isolation or in combination with BamHI. (One Hind III site is contributed by one of the primers and the other by the vector).

Two clones containing a D1.3 insert (fdTBam1) and fdTBam2) and one containing an NQ11 insert (NQ11Bam1)

were grown up and phage prepared as described earlier. ELISAs were carried out as described in example 6. No specific signal was found for any of these clones suggesting that the natural BamHI site is not a suitable site for insertion of a functional antibody (results not shown).

It may be possible to clone into alternative sites to retain binding activity. The peptide repeats present in gene III may provide such a site (figure 16 blocks A and B). This can be done by inserting a BamHI site and using the PCR product described above. To facilitate this, the natural BamHI site was removed by mutagenesis with the oligonucleotide G3mut6Bam shown below (using an in vitro mutagenesis kit (Amersham International)):-
 G3mut6Bam 5' CA AAC GAA TGG GTC CTC CTC ATT A 3'
 The underlined residue replaces an A residue, thereby removing the BamHI site. DNA was prepared from a number of clones and several mutants lacking BamHI sites identified by restriction digestion.

The oligonucleotide G3 Bamlink was designed to introduce a BamHI site at a number of possible sites within the peptide linker sites A and B, see figure 16(2). The sequence of the linker is:

Bamlink 5'CC (G or A) CC ACC CTC GGA TCC (G or A) CC ACC CTC 3'

Its relationship to the peptide repeats in gene III is shown in figure 16.

Example 14

PCR Assembly of Mouse VH and VL Kappa (VLK) Repertoires for Phage Display

The principle is illustrated in figure 17. Details are provided in sections A to F below but the broad outline is first discussed.

1. cDNA is prepared from spleen RNA from an appropriate mouse and the VH and VLK repertoires individually amplified. Separately, primers reverse and complementary to VH1FOR-2 (domain 1) and VLK2BACK (domain 2) are used to amplify an existing scFv-containing DNA by PCR. (The term FOR refers to e.g. a primer for amplification of sequences on the sense strand resulting in antisense coding sequences. The term BACK refers to e.g. a primer for amplification of sequences on the antisense strand resulting in sense coding sequences). This generates a 'linker' molecule encoding the linker with the amino acid sequence (1 letter code) (GGGGS)₃ which overlaps the two primary (VH and VLK) PCR products.
2. The separate amplified VH, VLK and linker sequences now have to be assembled into a continuous DNA molecule by use of an 'assembly' PCR. In the secondary 'assembly' PCR, the VH, VLK and linker bands are combined and assembled by virtue of the

above referred to overlaps. This generates an assembled DNA fragment that will direct the expression of VH and one VLK domain. The specific VH/VLK combination is derived randomly from the separate VH and VLK repertoires referred to above.

The assembly PCR is carried out in two stages. Firstly, 7 rounds of cycling with just the three bands present in the PCR, followed by a further 20 rounds in the presence of the flanking primers VH1BACK (referring to domain 1 of VH) and VLKFOR. The nucleotide sequences for these oligonucleotide primers are provided under the section entitled 'Primer Sequences' below. This two stage process, avoids the potential problem of preferential amplification of the first combinations to be assembled.

For cloning into the phage system, the assembled repertoires must be 'tagged' with the appropriate restriction sites. In the example provided below this is illustrated by providing an ApaI restriction site at the VH end of the continuous DNA molecule and a Not I site at the VLK end of the molecule. This is carried out by a third stage PCR using tagged primers. The nucleotide sequences for these oligonucleotide primers are also provided under the section entitled 'Primer Sequences' below. There are however, 4 possible kappa light chain sequences (whereas a single consensus heavy chain sequence can be used). Therefore 4 oligonucleotide primer sequences are provided for VLK.

For this third stage PCR, sets of primers which create the new restriction site and have a further 10 nucleotides on the 5' side of the restriction site have been used. However, long tags may give better cutting, in which case 15-20 nucleotide overhangs could be used.

Scrupulously clean procedures must be used at all times to avoid contamination during PCR. Negative controls containing no DNA must always be included to monitor for contamination. Gel boxes must be depurinated. A dedicated Geneclean kit (B10 101, Geneclean, La Jolla, San Diego, California, USA) can be used according to manufacturers instructions to extract DNA from an agarose gel. The beads, NaI and the NEW wash should be aliquoted.

All enzymes were obtained from CP Laboratories, P.O. Box 22, Bishop's Stortford, Herts CM20 3DH and the manufacturers recommended and supplied buffers were used unless otherwise stated.

A. RNA Preparation

RNA can be prepared using many procedures well known to those skilled in the art. As an example, the following protocol (Triton X-100 lysis, phenol/SDS RNase inactivation) gives excellent results with spleen and hybridoma cells (the addition of VRC (veronal ribosyl

complex) as an RNase inhibitor is necessary for spleen cells). Guanidinium isothiocyanate/CsCl procedures (yielding total cellular RNA) also give good results but are more time-consuming.

- 5 1. Harvest 1 to 5×10^7 cells by centrifugation in a bench top centrifuge at $800 \times g$ for 10 minutes at 4°C . Resuspend gently in 50ml of cold PBS buffer. Centrifuge the cells again at $800 \times g$ for 10 minutes at 4°C , and discard supernatant.
- 10 2. On ice, add 1 ml ice-cold lysis buffer to the pellet and resuspend it with a 1ml Gilson pipette by gently pipetting up and down. Leave on ice for 5 minutes.
3. After lysis, remove cell debris by centrifuging at 1300 rpm for 5 minutes in a microfuge at 4°C , in pre-cooled tubes.
- 15 4. Transfer 0.5 ml of the supernatant to each of two eppendorfs containing 60 μl 10% (w/v) SDS and 250 μl phenol (previously equilibrated with 100 mM Tris-HCl pH 8.0). Vortex hard for 2 minutes, then microfuge (13000 rpm) for five minutes at room temperature. Transfer the upper, aqueous, phase to a fresh tube.
- 20 5. Re-extract the aqueous upper phase five times with 0.5 ml of phenol.
6. Precipitate with 1/10 volume 3M sodium acetate and 2.5 volumes ethanol at 20°C overnight or dry ice-isopropanol for 30 minutes.
- 25 7. Wash the RNA pellet and resuspended in 50 μl to check concentration by OD260 and check 2 μg on a 1% agarose gel. 40 μg of RNA was obtained from spleen cells derived from mice.
- 30

Lysis buffer is [10mM Tris-HCl pH 7.4, 1mM MgCl_2 , 150mM NaCl, 10mM VRC (New England Biolabs), 0.5% (w/v) Triton X-100], prepared fresh.

- 35 Lysis buffer is [10mM Tris-HCl pH 7.4, 1mM MgCl_2 , 150mM NaCl, 10mM VRC (New England Biolabs), 0.5% (w/v) Triton X-100], prepared fresh.

B. cDNA Preparation

- 40 cDNA can be prepared using many procedures well known to those skilled in the art. As an example, the following protocol can be used:

1. Set up the following reverse transcription mix:

	μl
45 H_2O (DEPC-treated)	20
5mM dNTP	10
10 \times first strand buffer	10
0.1M DTT	10
FOR primer(s) (10 pmol/ μl)	2 (each) (see below)
RNasin (Promega; 40 U/ μl)	4

50 NB

- i) DEPC is diethylpyrocarbonate, the function of which is to inactivate any enzymes that could degrade DNA

or RNA

ii) dNTP is deoxynucleotide triphosphate
 iii) DTT is dithiothreitol the function of which is as an antioxidant to create the reducing environment necessary for enzyme function.

iv) RNasin is a ribonuclease inhibitor obtained from Promega Corporation, 2800 Woods Hollow Road, Madison, Wisconsin, USA.

2. Dilute 10 µg RNA to 40 µl final volume with DEPC-treated water. Heat at 65°C for 3 minutes and hold on ice for one minute (to remove secondary structure).

3. Add to the RNA the reverse transcription mix (58 µl) and 4 µl of the cloned reverse transcriptase 'Super RT' (Anglian Biotech Ltd., Whitehall House, Whitehall Road, Colchester, Essex) and incubate at 42°C for one hour.

4. Boil the reaction mix for three minutes, cool on ice for one minute and then spin in a microfuge to pellet debris. Transfer the supernatant to a new tube.

10 x first strand buffer is [1.4M KCl, 0.5M Tris-HCl pH 8.1 at 42°C 80mM MgCl₂].

The primers anneal to the 3' end. Examples of kappa light chain primers are MJK1FONX, MJK2FONX, MJK4FONX and MJK5FONX (provided under 'Primer Sequences' below) and examples of heavy chain primers are MIGG1, 2 (CTG GAC AGG GAT CCA GAG TTC CA) and MIGG3 (CTG GAC AGG GCT CCA TAG TTC CA) which anneal to CH1.

Alternatively, any primer that binds to the 3' end of the variable regions VH, VLK, VL, or to the constant regions CH1, CK or CL can be used.

C. Primary PCRs

For each PCR and negative control, the following reactions are set up (e.g. one reaction for each of the four VLKs and four VH PCRs). In the following, the Vent DNA polymerase sold by (C.P. Laboratories Ltd (New England Biolabs) address given above) was used. The buffers are as provided by C.P. Laboratories.

	µl
H ₂ O	32.5
10 x Vent buffer	5
20 x Vent BSA	2.5
5mM dNTPs	1.5
FOR primer 10 pmol/µl)	2.5
BACK primer 10pmol/µl	2.5

The FOR and BACK primers are given in the section below entitled 'Primer Sequences'. For VH, the FOR primer is VH1FOR-2 and the BACK primer is VH1BACK. For VLK the FOR primers are MJK1FONX, MJK2FONX, MJK4FONX and MJK5FONX (for the four respective kappa light chains) and the BACK

primer is VK2BACK. Only one kappa light chain BACK primer is necessary, because binding is to a nucleotide sequence common to the four kappa light chains.

5 UV this mix 5 minutes. Add 2.5 μ l cDNA preparation (from B above), 2 drops paraffin oil (Sigma Chemicals, Poole, Dorset, UK). Place on a cycling heating block, e.g. PHC-2 manufactured by Techne Ltd. Duxford UK, pre-set at 94°C. Add 1 μ l Vent DNA polymerase under the paraffin. Amplify using 25 cycles of 94°C 1 min, 72°C 2 min. Post-treat at 60°C for 5 min.

10 Purify on a 2% lmp (low melting point agarose/TAE (tris-acetate EDTA)gel and extract the DNA to 20 μ l H₂O per original PCR using a Geneclean kit (see earlier) in accordance with the manufacturers instructions.

15 D. Preparation of linker

Set up in bulk (e.g. 10 times)

		μ l
	H ₂ O	34.3
	10 x Vent buffer	5
20	20 x Vent BSA	2.5
	5mM dNTPs	2
	LINKFOR primer 10 pmol/ μ l)	2.5
	LINKBACK primer 10pmol/ μ l	2.5
	DNA from fcFv D1.3 (example 2)	1
25	Vent enzyme	0.2

The FOR and BACK primers are given in the section below entitled 'Primer Sequences'. The FOR primer is LINKFOR and the BACK primer is LINKBACK. Cover with paraffin and place on the cycling heating block (see above) at 94°C. Amplify using 25 cycles of 94°C 1 min, 65°C 1 min, 72°C 2 min. Post-treat at 60°C for 5 min.

30 Purify on 2% lmp/TAE gel (using a loading dye without bromophenol blue as a 93bp fragment is desired) and elute with SPIN-X column (Costar Limited, 205 Broadway, Cambridge, Ma. USA.,) and precipitation. Take up in 5 μ l H₂O per PCR reaction.

35 E. Assembly PCRs

A quarter of each PCR reaction product (5 μ l) is used for each assembly. The total volume is 25 μ l.

For each of the four VLK primers, the following are set up:

	H ₂ O	4.95
45	10 x Vent buffer	2.5
	20 x Vent BSA	1.25
	5mM dNTPs	0.8

50 UV irradiate this mix for 5 min. Add 5 μ l each of Vh and VK band from the primary PCRs and 1.5 μ l of linker as isolated from the preparative gels and extracted using the Geneclean kit as described in C and D above. Cover

with paraffin. Place on the cycling heating block preset at 94°C. Add 1ul Vent under the paraffin. Amplify using 7 cycles of 94°C 2 min, 72°C 4 min. Then return the temperature to 94°C.

- 5 Add 1.5ul each of VH1BACK and the appropriate VKFOR primers MJK1FONX, MJK2FONX, MJK4FONX or MJK5FONX (10 pmol/ul) at 94°C. The primers should have been UV-treated as above. Amplify using 20 cycles of 94°C 1.5 min, 72°C 2.5 min. Post-treat at 60°C for 5 min. Purify on 2% lmp/TAE gel and extract the DNA to 20ul H₂O per assembly PCR using a Geneclean kit (see earlier) in accordance with the manufacturers instructions.

F. Adding Restriction Sites

For each assembly and control set up:

15		<u>ul</u>
	H ₂ O	36.5
	10 x Taq buffer	5
	5mM dNTPs	2
	FOR primer (10 pmol/ul)	2.5
20	BACK primer (10 pmol/ul)	2.5
	Assembly product	1

- 25 The FOR and BACK primers are given in the section below entitled 'Primer Sequences'. The FOR primer is any of JK1NOT10, JK2NOT10, JK4NOT10 or JK5NOT10 (for the four respective kappa light chains) for putting a NotI restriction site at the VLK end. The BACK primer is HBKAPA10 for putting an ApaI restriction site at the VH end.

- 30 Cover with paraffin and place on the cycling heating block preset at 94°C. Add 0.5 ul Cetus Taq DNA polymerase (Cetus/perkin-Elmer, Beaconsfield, Bucks, UK) under the paraffin. Amplification is carried out using 11 to 15 rounds of cycling (depends on efficiency) at 94°C 1 min, 55°C 1 min, 72°C 2 min. Post-treat at 60°C for 5 min.

10 x Taq buffer is [0.1M Tris-HCl pH 8.3 at 25°C, 0.5M KCl, 15mM MgCl₂, 1mg/ml gelatin].

G. Work-up

- 40 Purify once with CHCl₃/IAA (isoamylalcohol), once with phenol, once with CHCl₃/IAA and back-extract everything to ensure minimal losses. Precipitate and wash twice in 70% EtOH. Dissolve in 70ul H₂O.

Digest overnight at 37°C with NotI:

45		<u>ul</u>
	DNA (joined seq)	70
	NEB NotI buffer x 10	10
	NEB BSA x 10	10
	NotI (10 U/ul)	10

- 50 The DNA (joined sequence) above refers to the assembled DNA sequence comprising in the 5' to 3' direction

 ApaI restriction site
 VH sequence

Linker sequence

VLK sequence

Not 1 restriction site.

5 The VLK sequence may be any one of four possible kappa chain sequences.

The enzymes Not 1 above, ApaI below and the buffers NEB Not 1, NEB BSA above and the NEB buffer 4 (below) are obtainable from CP Laboratories, New England Biolabs mentioned above.

10 Re-precipitate, take up in 80µl H₂O. Add to this 10µl NEB buffer 4 and 10µl ApaI 1.

Add the enzyme ApaI in aliquots throughout the day, as it has a short half-life at 37°C.

15 Purify on 2% Imp/TAE gel and extract the DNA using a Geneclean kit, in accordance with the manufacturers instructions. Redigest if desired.

H. Final DNA product

20 The final DNA product is an approximate 700 bp fragment with Apa I and NotI compatible ends consisting of randomly associated heavy and light chain sequences linked by a linker. A typical molecule of this type is the scFvD1.3 molecule incorporated into fdscFvD1.3 described in example 3. These molecules can then be ligated into suitable fd derived vectors, e.g. fdCAT2 (example 5), using standard techniques.

Primer sequences

Primary PCR oligos (restrictions sites underlined):

30 VH1FOR-2 TGA GGA GAC GGT GAC CGT GGT CCC TTG GCC CC
 VH1BACK AGG TSM ARC TGC AGS AGT CWG G
 MJK1FONX CCG TTT GAT TTC CAG CTT GGT GCC
 MJK2FONX CCG TTT TAT TTC CAG CTT GGT CCC
 MJK4FONX CCG TTT TAT TTC CAA CTT TGT CCC
 MJK5FONX CCG TTT CAG CTC CAG CTT GGT CCC
 VK2BACK GAC ATT GAG CTC ACC CAG TCT CCA
 35 Ambiguity codes M = A or C, R = A or G, S = G or C,
 W = A or T

PCR oligos to make linker:

40 LINKFOR TGG AGA CTC GGT GAG CTC AAT GTC
 LINKBACK GGG ACC ACG GTC ACC GTC TCC TCA

For adding restriction sites:

45 HBKAPA10 CAT GAC CAC AGT GCA CAG GTS MAR CTG CAG SAG TCW
 GG
 JKINOT10 GAG TCA TTC TGC GGC CGC CCG TTT GAT TTC CAG CTT
 GGT GCC
 JK2NOT10 GAG TCA TTC TGC GGC CGC CCG TTT TAT TTC CAG CTT
 GGT CCC
 JK4NOT10 GAG TCA TTC TGC GGC CGC CCG TTT TAT TTC CAA CTT
 TGT CCC
 50 JK5NOT10 GAG TCA TTC TGC GGC CGC CCG TTT CAG CTC CAG CTT
 GGT CCC

Example 15

Insertion of the Extracellular Domain of a Human Receptor for Platelet Derived Growth Factor (PDGF) isoform BB into fd CAT2

A gene fragment encoding the extracellular domain of the human receptor for platelet derived growth factor isoform BB (h-PDGFB-R) was isolated by amplification, using the polymerase chain reaction, of plasmid RP41, (from the American Type Culture collection, Cat. No.50735), a cDNA clone encoding amino-acids 43 to 925 of the PDGF-B receptor (Gronwald, R.G.K. et al PNAS 85 p3435-3439 (1988)). Amino acids 1 to 32 of h-PDGFB-R constitute the signal peptide. The oligonucleotide primers were designed to amplify the region of the h-PDGFB-R gene corresponding to amino acids 43 to 531 of the encoded protein. The primer RPDGF3 for the N-terminal region also included bases encoding amino acids 33 to 42 of the h-PDGFB-R protein (corresponding to the first ten amino acids from the N-terminus of the mature protein) to enable expression of the complete extracellular domain. The primers also incorporate a unique ApaI site at the N-terminal end of the fragment and a unique XhoI site at the C terminal end to facilitate cloning into the vector fdCAT2. The sequence of the primers is:

RPDGF3 5' CAC AGT GCA CTG GTC GTC ACA CCC CCG GGG CCA GAG
CTT GTC CTC AAT GTC TCC AGC ACC TTC GTT CTG 3'
RPDGF2 5' GAT CTC GAG CTT AAA GGG CAA GGA GTG TGG CAC 3'

PCR amplification was performed using high fidelity conditions (Eckert, K.A. and Kunkel, T.A. 1990 Nucl Acids Research 18 3739-3744). The PCR mixture contained: 20mM TrisHCl (pH7.3 at 70°C, 50mM KCl, 4mM magnesium chloride, 0.01% gelatin, 1mM each of dATP, dCTP, dGTP and dTTP, 500ng/ml RP41 DNA, 1µM each primer and 50 units/ml Taq polymerase (Cetus/Perkin Elmer, Beaconsfield, Bucks, U.K.). Thirty cycles of PCR were performed with denaturation at 92°C for 1 min, annealing at 60°C for 1min and extension at 72°C for 1.5 min. This reaction resulted in amplification of a fragment of ca. 1500bp as expected.

fdCAT2 vector DNA (see example 5) was digested with ApaI and XhoI (New England Biolabs) according to manufacturers recommendations, extracted with phenol/chloroform and ethanol precipitated (Sambrook et al, supra). Cloning of amplified RP41 DNA into this vector and identification of the desired clones was performed essentially as in example 7 except that digestion of the PCR product was with ApaI and Xho I. Colonies containing h-PDGFB-R DNA were identified by probing with 32p labelled RPDGF2 and the presence of an insert in hybridising colonies was confirmed by analytical PCR using RPDGF3 and RPDGF2 using the

conditions described in example 7.

Example 16

5 Binding of ^{125}I -PDGF-BB to the Extracellular Domain of
the Human Receptor for Platelet Derived Growth Factor
Isoform BB Displayed on the Surface of fd Phage.
Measured using an Immunoprecipitation Assay.

10 Phage particles, expressing the extracellular domain
 of the human platelet derived growth factor isoform BB
 receptor (fd h-PDGFB-R), were prepared by growing E.coli
 MC1061 cells transformed with fd h-PDGFB-R in 50ml of
 2xTY medium with 15ug/ml tetracycline for 16 to 20 hours.
 15 Phage particles were concentrated using polyethylene
 glycol as described in example 6 and resuspended in PDGF
 binding buffer (25mM HEPES, pH7.4, 0.15mM NaCl, 1mM
 magnesium chloride, 0.25% BSA) to 1/33rd of the original
 volume. Residual bacteria and undissolved material were
 removed by spinning for 2 min in a microcentrifuge.
 20 Immunoblots using an antiserum raised against gene III
 protein (Prof. I. Rashed, Konstanz, Germany) show the
 presence in such phage preparations of a geneIII-h-PDGFB-
 R protein of molecular mass 125000 corresponding to a
 fusion between h-PDGFB-R external domain (55000 daltons)
 and geneIII (apparent molecular mass 70000 on SDS-
 25 polyacrylamide gel).

Duplicate samples of 35ul concentrated phage were
 incubated with ^{125}I -PDGF-BB (78.7fmol, 70nCi, 882Ci/mmol;
 Amersham International plc, Amersham, Bucks) for 1 hour
 at 37°C. Controls were included in which fdTPs/Bs vector
 30 phage (figure 4) or no phage replaced fd h-PDGFB-R phage.
 After this incubation, 10ul of sheep anti-M13 polyclonal
 antiserum (a gift from M. Hobart) was added and
 incubation continued for 30 min at 20°C. To each sample,
 40ul (20ul packed volume) of protein G Sepharose Fast
 35 Flow (Pharmacia, Milton Keynes) equilibrated in PDGF
 binding buffer was added. Incubation was continued for
 30 min at 20°C with mixing by end over end inversion on a
 rotating mixer. The affinity matrix was spun down in a
 microcentrifuge for 2 min and the supernatant removed by
 40 aspiration. Non-specifically bound ^{125}I -PDGF-BB was
 removed by resuspension of the pellet in 0.5ml PDGF
 binding buffer, mixing by rotation for 5 min,
 centrifugation and aspiration of the supernatant,
 followed by two further washes with 0.5ml 0.1% BSA, 0.2%
 45 Triton-X-100. The pellet finally obtained was
 resuspended in 100ul PDGF binding buffer and counted in a
 Packard gamma counter. For displacement studies,
 unlabelled PDGF-BB (Amersham International) was added to
 the stated concentration for the incubation of ^{125}I -PDGF-
 50 BB with phage.

^{125}I -PDGF-BB bound to the fd h-PDGFB-R phage and was
 immunoprecipitated in this assay. Specific binding to

receptor phage was 3.5 to 4 times higher than the non-specific binding with vector phage fdTPs/Bs or no phage (fig. 19). This binding of ^{125}I -PDGF-BB could be displaced by the inclusion of unlabelled PDGF-BB in the incubation with phage at 37°C (fig. 20). At 50nM, unlabelled PDGF-BB the binding of ^{125}I -PDGF-BB was reduced to the same level as the fdTPs/Bs and no phage control. Figure 21 shows the same data, but with the non-specific binding to vector deducted.

These results indicate that a specific saturable site for ^{125}I -PDGF-BB is expressed on fd phage containing cloned h-PDGFB-R DNA. Thus, the phage can display the functional extracellular domain of a cell surface receptor.

Example 17, Construction of Phagemid Containing GeneIII fused with the Coding Sequence for a Binding Molecule

It would be useful to improve the transfection efficiency of the phage-binding molecule system and also to have the possibility of displaying different numbers and specificities of binding molecules on the surface of the same bacteriophage. The applicants have devised a method that achieves both aims.

The approach is derived from the phagemid system based on pUC119 [Vieira, J and Messing, J. (1987) Methods Enzymol. 153:3]. In brief, gene III from fd-CAT2 (example 5) and gene III scFv fusion from fd-CAT2 scFv D1.3 (example 2) were cloned downstream of the lac promoter in separate samples of pUC119, in order that the inserted gene III and gene III fusion could be 'rescued' by M13M07 helper phage [Vieira, J and Messing, J. et supra.] prepared according to Sambrook et al. 1989 supra. The majority of rescued phage would be expected to contain a genome derived from the pUC119 plasmid that contains the binding molecule-gene III fusion and should express varying numbers of the binding molecule on the surface up to the normal maximum of 3-5 molecules of gene III of the surface of wild type phage. The system has been exemplified below using an antibody as the binding molecule.

An fdCAT2 containing the single chain Fv form of the D1.3 antilysozyme antibody was formed by digesting fdTscFvD1.3 (example 2) with PstI and XhoI, purifying the fragment containing the scFv fragment and ligating this into PstI and XhoI digested fdCAT2. The appropriate clone, called fdCAT2 scFvD1.3 was selected after plating onto 2xTY tetracycline (15µg/ml) and confirmed by restriction enzyme and sequence analysis.

Gene III from fd-CAT2 (example 5) and the gene III scFv fusion from fd-CAT2 scFvD1.3 was PCR-amplified using the primers A and B shown below:

Primer A: TGC GAA GCT TTG GAG CCT TTT TTT TTG GAG ATT TTC
AAC G

Primer B: CAG TGA ATT CCT ATT AAG ACT CCT TAT TAC GCA GTA
TGT TAG C

Primer A anneals to the 5' end of gene III including the ribosome binding site is located and incorporates a Hind III site. Primer B anneals to the 3' end of gene III at the C-terminus and incorporates two UAA stop codons and an EcoRI site. 100 ng of fd-CAT2 and fd-CAT2 scFv D1.3 DNA was used as templates for PCR-amplification in a total reaction volume of 50µl as described in example 7, except that 20 cycles of amplification were performed: 94°C 1 minute, 50°C 1 minute, 72°C 3 minutes. This resulted in amplification of the expected 1.2Kb fragment from fd-CAT2 and a 1.8Kb fragment from fd-CAT2 scFv D1.3.

The PCR fragments were digested with EcoRI and Hind III, gel-purified and ligated into Eco-R1- and Hind III-cut and dephosphorylated pUC119 DNA and transformed into E.coli TG1 using standard techniques (Sambrook et al., et supra). Transformed cells were plated on SOB agar (Sambrook et al. 1989 supra) containing 100µg/ml ampicillin and 2% glucose. The resulting clones were called pCAT-3 (derived from fd-CAT2) and pCAT-3 scFv D1.3 (derived from fd-CAT2 scFv D1.3).

Example 18. Rescue of Anti-Lysozyme Antibody Specificity from pCAT-3 scFv D1.3 by M13K07

Single pCAT-3 and pCAT-3 scFv D1.3 colonies were picked into 1.5ml 2TY containing 100µg/ml ampicillin and 2% glucose, and grown 6 hrs at 30°C. 30µl of these stationary cells were added to 6mls 2TY containing 100µg/ml ampicillin and 2% glucose in 50ml polypropylene tubes (Falcon, Becton Dickinson Labware, 1950 Williams Drive, Oxnard, CA. USA) and grown for 1.5 hrs at 30°C at 380rpm in a New Brunswick Orbital Shaker (New Brunswick Scientific Ltd., Edison House 163 Dixons Hill road, North Mimms, Hatfield, UK). Cells were pelleted by centrifugation at 5,000g for 25 minutes and the tubes drained on tissue paper. The cell pellets were then suspended in 6mls 2TY containing 1.25×10^9 p.f.u. ml⁻¹ M13K07 bacteriophage added. The mixture was left on ice for 5 minutes followed by growth at 35°C for 45 minutes at 450rpm. A cocktail was then added containing 4µl 100µg/ml ampicillin, 0.5µl 0.1M IPTG and 50µl 10mg/ml kanamycin, and the cultures grown overnight at 35°C, 450rpm.

The following day the cultures were centrifuged and phage particles PEG precipitated as described in example 6. Phage pellets were resuspended in 100µl TE (tris-EDTA see example 6) and phage titred on E.coli TG1. Aliquots of infected cells were plated on 2TY containing either 100µg/ml ampicillin to select for pUC119 phage particles, or 50µg/ml kanamycin to select for the M13 K07 helper phage. Plates were incubated overnight at 37°C and

antibiotic-resistant colonies counted:

DNA	amp ^R	kan ^R
pCAT-3	1.8x10 ¹¹ colonies	1.2x10 ⁹ colonies
pCAT-3scFv D1.3	2.4x10 ¹¹ colonies	2.0x10 ⁹ colonies

5 This shows that the amp^R phagemid particles are infective and present in the rescued phage population at a 100-fold excess over kan^R M13K07 helper phage.

10 Phage were assayed for anti-lysozyme activity by ELISA as described in example 6, with the following modifications:

1) ELISA plates were blocked for 3 hrs with 2% Marvel/PBS.

15 2) 50µl phage, 400µl 1xPBS and 50µl 20% Marvel were mixed end over end for 20 minutes at room temperature before adding 150µl per well.

3) Phage were left to bind for 2 hours at room temperature.

4) All washes post phage binding were:

20 2 quick rinses PBS/0.5% Tween 20

3x2 minute washes PBS/0.5% Tween 20

2 quick rinses PBS no detergent

3x2 minute washes PBS no detergent

25 The result of this ELISA is shown in figure 22, which shows that the antibody specificity can indeed be rescued efficiently.

30 It is considered a truism of bacterial genetics that when mutant and wild-type proteins are co-expressed in the same cell, the wild-type proteins are co-expressed in same cell, the wild-type protein is used preferentially. This is analogous to the above situation wherein mutant (i.e. antibody fusion) and wild-type gene III proteins (from M13K07) are competing for assembly as part of the pUC119 phagemid particle. It is therefore envisaged that the majority of the resulting pUC 119 phage particles will have fewer gene III-antibody fusion molecules on their surface than is the case for purely phage system described for instance in example 2. Such phagemid antibodies are therefore likely to bind antigen with a lower avidity than fd phage antibodies with three or more copies of the antibody fusion on their surfaces (there is no wild-type gene III, in the system described, for instance, in example 2), and provide a route to production of phage particles with different numbers of the same binding molecule (and hence different acidities for the ligand/antigen) or multiple different binding specificities on their surface, by using helper phage such as M13K07 to rescue cells expressing two or more gene III-antibody fusions.

50 It is also possible to derive helper phage that do not encode a functional gene III in their genomes (by for example deleting the gene III sequence or a portion of it or by incorporating an amber mutation within the gene).

These defective phages will only grow on appropriate cells (for example that provide functional gene III in trans, or contain an amber supressor gene), but when used to rescue phage antibodies, will only incorporate the gene III antibody fusion encoded by the phagemid into the released phage particle.

Example 19. Transformation Efficiency of pCAT-3 and pCAT-3 scFv D1.3 phagemids

pUC 19, pCAT-3 and pCAT-3 scFv D1.3 plasmid DNAs, and fdCAT-2 phage DNA was prepared, and used to transform E.coli TG1, pCAT-3 and pCAT-3 scFv D1.3 transformations were plated on SOB agar containing 100µg/ml ampicillin and 2% glucose, and incubated overnight at 30°C. fdCAT-2 transformations were plated on TY agar containing 15µg/ml tetracycline and incubated overnight at 37°C. Transformation efficiencies are expressed as colonies per µg of input DNA.

DNA	Transformation efficiency
pUC 19	1.10^9
pCAT-3	1.10^8
pCAT-3scFv D1.3	1.10^8
fd CAT-2	8.10^5

As expected, transformation of the phagemid vector is approximately 100-fold more efficient than the parental fdCAT-2 vector. Furthermore, the presence of a scFv antibody fragment does not compromise efficiency. This improvement in transformation efficiency is practically useful in the generation of phage antibody libraries that have large repertoires of different binding specificities.

Example 20

PCR Assembly of a Single Chain Fv Library from an Immunised Mouse

To demonstrate the utility of phage for the selection of antibodies from repertoires, the first requirement is to be able to prepare a diverse, representative library of the antibody repertoire of an animal and display this repertoire on the surface of bacteriophage fd.

Cytoplasmic RNA was isolated according to example 14 from the pooled spleens of five male Balb/c mice boosted 8 weeks after primary immunisation with 2-phenyl-5-oxazolone (ph OX) coupled to chicken serum albumin. cDNA preparation and PCR assembly of the mouse VH and VL kappa repertoires for phage display was as described in example 14. The molecules thus obtained were ligated into fdCAT2.

Vector fdCAT2 was extensively digested with NotI and ApaLI., purified by electroelution (Sambrook et al. 1989 supra) and 1 µg ligated to 0.5 µg (5 µg for the hierarchical libraries: see example 22) of the assembled scFv genes in 1 ml with 8000 units T4 DNA ligase (New

England Biolabs). The ligation was carried out overnight at 16°C. Purified ligation mix was electroporated in six aliquots into MC1061 cells (W. J. Dower, J. F. Miller & C. W. Ragsdale Nucleic Acids Res. 16 6127-6145 1988) and

5 plated on NZY medium (Sambrook et al. 1989 supra) with 15µg/ml tetracycline, in 243x243 mm dishes (Nunc): 90-95% of clones contained scFv genes by PCR screening. Recombinant colonies were screened by PCR (conditions as in example 7 using primers VH1BACK and MJK1FONX, 10 MJK2FONX, MJK4FONX and MJK5FONX (see example 14) followed by digestion with the frequent cutting enzyme BstNI (New England Biolabs, used according to the manufacturers instructions). The library of 2×10^5 clones appeared 15 diverse as judged by the variety of digestion patterns seen in Figure 23, and sequencing revealed the presence of most VH groups (R. Dildrop, Immunol. Today 5 85-86. 1984) and VK subgroups (Kabat. E.A. et al. 1987 supra) (data not shown). None of the 568 clones tested bound to phOx as detected by ELISA as in example 9.

20 Thus the ability to select antibody provided by the use of phage antibodies (as in example 21) is essential to readily isolate antibodies with antigen binding activity from randomly combined VH and VL domains. Very 25 extensive screening would be required to isolate antigen-binding fragments if the random combinatorial approach of Huse et al. 1989 (supra) were used.

Example 21

Selection of Antibodies Specific for 2-phenyl-5-oxazolone from a Repertoire Derived from an Immunised Mouse

30 The library prepared in example 20 was used to demonstrate that ability of the phage system to select antibodies on the basis of their antibody specificity.

None of the 568 clones tested from the unselected library bound to phOx as detected by ELISA.

35 Screening for binding of the phage to hapten was carried out by ELISA: 96-well plates were coated with 10 µg/ml phOx-BSA or 10 µg/ml BSA in phosphate-buffered saline (PBS) overnight at room temperature. Colonies of phage-transduced bacteria were inoculated into 200 µl 2 x 40 TY with 12.5 µg/ml tetracycline in 96-well plates ('cell wells', Nuclon) and grown with shaking (300 rpm) for 24 hours at 37°C. At this stage cultures were saturated and phage titres were reproducible (10^{10} TU/ml). 50 µl phage supernatant, mixed with 50 µl PBS containing 4% skimmed 45 milk powder, was then added to the coated plates. Further details as in example 9.

The library of phages was passed down a phOx affinity column (Table 4A), and eluted with hapten. Colonies from the library prepared in example 22 were 50 scraped into 50ml 2 x TY medium³⁷ and shaken at 37°C for 30 min. Liberated phage were precipitated twice with polyethylene glycol and resuspended to 10^{12} TU

(transducing units)/ml in water (titred as in example 8). For affinity selection, a 1 ml column of phOx-BSA-Sepharose (O. Makela, M. Kaartinen, J.L.T. Pelonen and K. Karjalainen J. Exp. Med. 148 1644-1660, 1978) was washed with 300 ml phosphate-buffered saline (PBS), and 20 ml PBS containing 2% skimmed milk powder (MPBS). 10¹² TU phage were loaded in 10 ml MPBS, washed with 10 ml MPBS and finally 200 ml PBS. The bound phage were eluted with 5 ml 1 mM 4- ϵ -amino-caproic acid methylene 2-phenyl-oxazol-5-one (phOx-CAP; O. Makela et al. 1978, supra). About 10⁶ TU eluted phage were amplified by infecting 1 ml log phase E.coli TG1 and plating as above. For a further round of selection, colonies were scraped into 10 ml 2 x TY medium and then processed as above. Of the eluted clones, 13% were found to bind to phOx after the first round selection, and ranged from poor to strong binding in ELISA.

To sequence clones, template DNA was prepared from the supernatants of 10 ml cultures grown for 24 hours, and sequenced using the dideoxy method and a Sequenase kit (USB), with primer LINKFOR (see example 14) for the VH genes and primer fdSEQ1 (5'-GAA TTT TCT GTA TGA GG) for the Vk genes. Twenty-three of these hapten-binding clones were sequenced and eight different VH genes (A to H) were found in a variety of pairings with seven different Vk genes (a to g) (Fig. 24). Most of the domains, such as VH-B and Vk-d were 'promiscuous', able to bind hapten with any of several partners.

The sequences of the V-genes were related to those seen in the secondary response to phOx, but with differences (Fig. 24). Thus phOx hybridomas from the secondary response employ somatically mutated derivatives of three types of Vk genes - Vkox1. 'Vkox-like' and Vk45.1 genes (C. Berek, G. M. Griffiths & C. Milstein Nature 316 412-418 (1985)). These can pair with VH genes from several groups, from Vkox1 more commonly pairs with the VHox1 gene (VH group 2. R.Dildrop uupra). Vkox1 genes are always, and Vkox-like genes often, found in association with heavy chains (including VHox1) and contain a short five residue CDR3, with the sequence motif Asp-X-Gly-X-X in which the central glycine is needed to create a cavity for phOx. In the random combinatorial library however, nearly all of the VH genes belonged to group 1, and most of the Vk genes were ox-like and associated with VH domains with a five residue CDR3, motif Asp/Asn-X-Gly-X-X (Fig. 24). Vkox1 and VHox1 were found only once (Vk-f and VH-E), and not in combination with each other. Indeed Vk-f lacks the Trp91 involved in phOx binding and was paired with a VH (VH-C) with a six residue CDR3.

A matrix combination of VH and VK genes was identified in phOx-binding clones selected from this

random combinational library. The number of clones found with each combination are shown in Fig. 25. The binding to phOx-BSA, as judged by the ELISA signal, appeared to vary (marked by shading in Fig. 25). No binding was seen to BSA alone.

A second round of selection of the original, random combinational library from immune mice resulted in 93% of eluted clones binding phOx (Table 4). Most of these clones were Vk-d combinations, and bound strongly to phOx in ELISA (data not shown). Few weak binders were seen. This suggested that affinity chromatography had not only enriched for binders, but also for the best.

Florescence quench titrations determined the Kd of VH-B/Vk-d for phOx-GABA as 10^{-8} M (example 23), indicating that antibodies with affinities representative of the secondary response can be selected from secondary response, only two (out of eleven characterised) secrete antibodies of a higher affinity than VH-B/Vk-d (C. Berek et al. 1985 supra). The Kd of VH-B/Vk-b for phOx-GABA was determined as 10^{-5} M (example 23). Thus phage bearing scFv fragments with weak affinities can be selected with antigen, probably due to the avidity of the multiple antibody heads on the phage.

This example shows that antigen specificities can be isolated from libraries derived from immunised mice. It will often be desired to express these antibodies in a soluble form for further study and for use in therapeutic and diagnostic applications. Example 23 demonstrates determination of the affinity of soluble scFv fragments selected using phage antibodies. Example 27 demonstrates that soluble fragments have similar properties to those displayed on phage. For many purposes it will be desired to construct and express an antibody molecule which contains the Fc portions of the heavy chain, and perhaps vary the immunoglobulin isotype. To accomplish this, it is necessary to subclone the antigen binding sites identified using the phage selection system into a vector for expression in mammalian cells, using methodology similar to that described by Orlandi, R. et al. (1989, supra). For instance, the VH and VL genes could be amplified separately by PCR with primers containing appropriate restriction sites and inserted into vectors such as pSV-gpt HuIgG1 (L. Riechmann et al Nature 332 323-327), 1988) which allows expression of the VH domain as part of a heavy chain IgG1 isotype and pSV-hyg HuCK which allows expression of the VL domain attached to the K light chain constant region. Furthermore, fusions of VH and VL domains can be made with genes encoding non-immunoglobulin proteins, for example, enzymes.

Example 22

Generation of Further Antibody Specificities by the Assembly of Hierarchical Libraries

Further antibody specificities were derived from the library prepared and screened in examples 20 and 21 using a hierarchical approach.

5 The promiscuity of the VH-B and Vk-d domains prompted the applicants to force further pairings, by assembling these genes with the entire repertoires of either Vk or VH genes from the same immunised mice. The resulting 'hierarchical' libraries, (VH-B x Vk-rep and VH-rep x Vk-d), each with 4×10^7 members, were subjected to a round of selection and hapten-binding clones isolated (Table 4). As shown by ELISA, most were strong binders. By sequencing twenty-four clones from each library, the applicants identified fourteen new partners for VH-B and thirteen for Vk-d (Fig. 24). Apart from VH-B and Vk-c, none of the previous partners (or indeed other clones) from the random combinatorial library was isolated again. Again the Vk genes were mainly ox-like and the VH genes mainly group 1 (as defined in Dildrop, R. 1984 supra), but the only examples of Vkoxl (Vk-h, -p, -q and -r) have Trp91, and the VH-CDR3 motif Asp-X-Gly-X-X now predominates. Thus some features of the phoX hybridomas seemed to emerge more strongly in the hierarchical library. The new partners differed from each other mainly by small alterations in the CDRs, indicating that much of the subtle diversity had remained untapped by the random combinatorial approach. More generally it has been shown that a spectrum of related antibodies can be made by keeping one of the partners fixed and varying the other, and this could prove invaluable for fine tuning of antibody affinity and specificity.

Therefore, again, phage antibodies allow a greater range of antibody molecules to be analysed for desired properties.

35 This example, and example 21, demonstrate the isolation of individual antibody specificities through display on the surface of phage. However, for some purposes it may be more desirable to have a mixture of antibodies, equivalent to a polyclonal antiserum (for instance, for immunoprecipitation). To prepare a mixture of antibodies, one could mix clones and express soluble antibodies or antibody fragments or alternatively select clones from a library to give a highly enriched pool of genes encoding antibodies or antibody fragments directed against a ligand of interest and express antibodies from these clones.

45 Example 23

Selection of Antibodies Displayed on Bacteriophage with Different Affinities for 2-phenyl-5-oxazolone using Affinity Chromatography

50 The ELISA data shown in example 21 suggested that affinity chromatography had not only enriched for binders, but also for the best. To confirm this, the

binding affinities of a strong binding and a weak binding phage were determined and then demonstrated that they could be separated from each other using affinity chromatography.

- 5 Clones VH-B/Vk-b and VH-B/Vk-d were reamplified with MJK1FONX, MJK2FONX, MJK4FONX and MJK5FONX (see example 14) and VH1BACK-SfiI (5'-TCG CGG CCC AGC CGG CCA TGG CC(G/C) AGG T(C/G)(A/C) A(A/G)C TGC AG(C/G) AGT C(A/T)G G), a primer that introduces an SfiI site (underlined) at the 5' end of the VH gene. VH-B/Vk-d was cloned into a phagemid e.g. pJM1 (a gift from A. Griffiths and J. Marks) as an SfiI-NotI cassette, downstream of the pelB leader for periplasmic secretion (M. Better et al. supra), with a C-terminal peptide tag for detection (see example 24 and figure), and under the control of a P_L promoter (H. Shimatake & M. Rosenberg Nature 292 128-132 1981). The phagemid should have the following features: a) unique SfiI and NotI restriction sites downstream of a pelB leader; b) a sequence encoding a C-terminal peptide tag for detection; and c) a P_L promoter controlling expression. 10 litre cultures of E.coli N4830-1 (M. E. Gottesman, S. Adhya & A. Das J.Mol.Biol 140 57-75 1980) harbouring each phagemid were induced as in K. Nagai & H. C. Thogerson (Methods Enzymol 153 461-481 1987) and supernatants precipitated with 50% ammonium sulphate. The resuspended precipitate was dialysed into PBS + 0.2 mM EDTA (PBSE), loaded onto a 1.5ml column of phOx:Sephacrose and the column washed sequentially with 100 ml PBS: 100 ml 0.1 M Tris-HCl, 0.5 M NaCl, pH 8.0: 10ml 50 mM citrate, pH 5.0: 10 ml 50 mM citrate, pH4.0, and 20 ml 50 mM glycine, pH 3.0. scFv fragments were eluted with 50 mM glycine, pH 2.0, neutralised with Tris base and dialysed against PBSE. VH-B/Vk-b was cloned into a phagemid vector based on pUC119 encoding identical signal and tag sequences to pJM1, and expression induced at 30°C in a 10 litre culture of E.coli TG1 harbouring the phagemid as in D. de Bellis & I. Schwartz (1980 Nucleic Acids Res 18 1311). The low affinity of clone VH-B/Vk-b made its purification on phOx-Sephacrose impossible. Therefore after concentration by ultrafiltration (Filtron, Flowgen), the supernatant (100 ml of 600 ml) was loaded onto a 1 ml column of protein A-Sephacrose coupled (E. Harlow & D. Lane 1988 supra) to the monoclonal antibody 9E10 (Evan, G. I. et al. Mol.Cell Biol. 5 3610-3616 1985) that recognises the peptide tag. The column was washed with 200 ml PBS and 50 ml PBS made 0.5 M in NaCl. scFv fragments were eluted with 100 ml 0.2M glycine, pH 3.0, with neutralisation and dialysis as before.
- 50 The K_d ($1.0 \pm 0.2 \times 10^{-8}$ M) for clone VH-B/Vk-d was determined by fluorescence quench titration with 4-E-amino-butyrilic acid methylene 2-phenyl-oxazol-5-one (phOx-

GABA Co. Makela et al, 1978 supra). Excitation was at 280 nm, emission was monitored at 340 nm and the K_d calculated. The K_d of the low affinity clone VH-B/Vk-b was determined as $1.8 \pm 0.3 \times 10^{-5}$ M (not shown). To minimise light adsorption by the higher concentrations of phOx-GABA required, excitation was at 260 nm and emission was monitored at 304 nm. In addition the fluorescence values were divided by those from a parallel titration of the lysozyme binding Fv fragment D1.3. The value was calculated as in H. N. Eisen Meth.Med.Res. 10 115-121 1964. A mixture of clones VH-B/Vk-b and VH-B/Vk-d, 7×10^{10} TU phage in the ratio 20 VH-B/Vk-b : 1 VH-B/Vk-d were loaded onto a phOx-BSA-Sepharose column in 10 ml MPBS and eluted as above. Eluted phage were used to reinfect E.coli TGI, and phage produced and harvested as before. Approximately 10^{11} TU phage were loaded onto a second affinity column and the process repeated to give a total of three column passes. Dilutions of eluted phage at each stage were plated in duplicate and probed separately with oligonucleotides specific for Vk-b (5'GAG CGG GTA ACC ACT GTA CT) or Vk-d (5'-GAA TGG TAT AGT ACT ACC CT). After these two rounds, essentially all the eluted phage were VH-B/Vk-d (table 4). Therefore phage antibodies can be selected on the basis of the antigen affinity of the antibody displayed.

Example 24

Construction of Phagemid pHEN1 for the Expression of Antibody Fragments Expressed on the Surface of Bacteriophage following Superinfection

The phagemid pHEN1 (figure 26) is a derivative of pUC119 (Vieira, J. & Messing, J. Methods Enzymol 153 pp 3-11, 1987). The coding region of g3p from fdCAT2, including signal peptide and cloning sites, was amplified by PCR, using primers G3FUFO and G3FUBA (given below) (which contain EcoRI and HindIII sites respectively), and cloned as a HindIII-EcoRI fragment into pUC119. The HindIII-NotI fragment encoding the g3p signal sequence was the replaced by a pelB signal peptide (Better, M. et al. Science 240 1041-1043, 1988) with an internal SfiI site, allowing antibody genes to be cloned as fII-NotI fragments. A peptide tag, c-myc, (Munro, S. & Pelham, H. Cell 46 291-300, 1986) was introduced directly after the NotI site by cloning an oligonucleotide cassette, and followed by an amber codon introduced by site-directed mutagenesis using an in vitro mutagenesis kit (Amersham International) (figure 26b).

G3FUFO, 5'-CAG TGA ATT CTT ATT AAG ACT CCT TAT TAC GCA GTA
TGT TAG C;
G3FUBA, 5'-TGC GAA GCT TTG GAG CCT TTT TTT TTG GAG ATT TTC
AAC G;

Example 25

Display of Single Chain Fv and Fab Fragments Derived from the Anti-Oxazolone Antibody NQ10.12.5 on Bacteriophage fd using pHEN1 and fdCAT2

A range of constructs (see figure 27) were made from a clone (essentially construct II in pUC19) designed for expression in bacteria of a soluble Fab fragment (Better et al. 1988 see above) from the mouse anti-phOx (2-phenyl-5-oxazolone) antibody NQ10.12.5 (Griffiths, G. M. et al. Nature 312, 271-275, 1984). In construct II, the V-regions are derived from NQ10.12.5 and attached to human C κ and C μ 1 (γ 1 isotype) constant domains. The C-terminal cysteine residues, which normally form a covalent link between light and heavy antibody chains, have been deleted from both the constant domains. To clone heavy and light chain genes together as Fab fragments (construct II) or as separate chains (constructs III and IV) for phage display, DNA was amplified from construct II by PCR to introduce a NotI restriction site at the 3' end, and at the 5' end either an ApaLI site (for cloning into fd-CAT2) or SfiI site (for cloning into pHEN1). The primers FABNOTFOK with VH1BACKAPA (or VH1BACKSFI15) were used for PCR amplification of genes encoding Fab fragments (construct II), the primers FABNOTFOH with VH1BACKAPA (or VH1BACKSFI15) for heavy chains (construct III), and the primers FABNOTFOK and MVKBAAPA (or MVKBASFI) for light chains (construct IV).

The single-chain Fv version of NQ10.12.5 (construct I) has the heavy (VH) and light chain (VK) variable domains joined by a flexible linker (Gly₄Ser)₃ (Huston, J. S. et al. Proc. Natl. Acad. Sci. USA 85 5879-5883, 1988) and was constructed from construct II by 'splicing by overlap extension' as in example 14. The assembled genes were reamplified with primers VK3F2NOT and VH1BACKAPA (or VH1BACKSFI15) to append restriction sites for cloning into fd-CAT2 (ApaLI-NotI) or pHEN1 (SfiI-NotI).

VH1BACKAPA, 5'-CAT GAC CAC AGT GCA CAG GT(C/G) (A/C)A(A/G) CTG CAG (C/G)AG TC(A/T) GG;
VH1BACKSFI15, 5'-CAT GCC ATG ACT CGC GGC CCA GCC GGC CAT GGC C(C/G)A GGT (C/G)(A/C)A (A/G)CT GCA G(C/G)A GTC (A/T)GG;
FABNOTFOH, 5'-CCA CGA TTC TGC GGC CGC TGA AGA TTT GGG CTC AAC TTT CTT GTC GAC;
FABNOTFOK, 5'-CCA CGA TTC TGC GGC CGC TGA CTC TCC GCG GTT GAA GCT CTT TGT GAC;
MVKBAAPA, 5'-CAC AGT GCA CTC GAC ATT GAG CTC ACC CAG TCT CCA;
MVKBASFI, 5'-CAT GAC CAC GCG GCC CAG CCG GCC ATG GCC GAC

ATT GAG CTC ACC CAG TCT CCA;
 VK3F2NOT, 5'-TTC TGC GGC CGC CCG TTT CAG CTC GAG CTT GGT
 CCC.

Restriction sites are underlined.

5 Rescue of Phage and Phagemid particles

Constructs I-IV (figure 27) were introduced into both fd-CAT2 and pHEN1. Phage fd-CAT2 (and fd-CAT2-I,II,III or IV) was taken from the supernatant of infected E.coli TG1 after shaking at 37°C overnight in 2xTY medium with 12.5µg/ml tetracycline, and used directly in ELISA. Phagemid pHEN1 (and pHEN1-I and II) in E.coli TG1 (supE) were grown overnight in 2 ml 2xTY medium, 100 µg/ml ampicillin, and 1% glucose (without glucose, expression of g3p prevents later superinfection by helper phage). 10µl of the overnight culture was used to inoculate 2 ml of 2xTY medium, 100µg/ml ampicillin, 1% glucose, and shaken at 37°C for 1 hour. The cells were washed and resuspended in 2xTY, 100 µg/ml ampicillin, and aphagemid particles rescued by adding 2 µl (10⁸pfu) VCSM13 helper phage (Stratagene). After growth for one hour, 4µl kanamycin (25 mg/ml) was added, and the culture grown overnight. The phagemid particles were concentrated 10-fold for ELISA by precipitation with polyethylene glycol.

ELISA
 25 Detection of phage binding to 2-phenyl-5-oxazolone (phOx) was performed as in example 9. 96-well plates were coated with 10 µg/ml phOx-BSA or 10 µg/ml BSA in PBS overnight at room temperature, and blocked with PBSS containing 2% skimmed milk powder. Phage (mid) supernatant (50 µl) mixed with 50 µl PBS containing 4% skimmed milk powder was added to the wells and assayed. To detect binding of soluble scFv or Fab fragments secreted from pHEN1, the c-myc peptide tag described by Munro and Pelham 1986 supra, was detected using the anti-myc monoclonal 9E10 (Evan, G. I. et al. Mol Cell Biol 5 3610-3616, 1985) followed by detection with peroxidase-conjugated goat anti-mouse immunoglobulin. Other details are as in example 9.

The constructs in fdCAT2 and pHEN1 display antibody fragments of the surface of filamentous phage. The phage vector, fd-CAT2 (figure 8) is based on the vector fd-tet (Zacher, A. N. et al. Gene 9 127-140, 1980) and has restriction sites (ApaI and NotI) for cloning antibody genes (or other protein) genes for expression as fusions to the N-terminus of the phage coat protein g3p. Transcription of the antibody-g3p fusions in fd-CAT2 is driven from the gene III promoter and the fusion protein targetted to the periplasm by means of the g3p leader. Fab abd scFv fragments of NQ10.12.5 cloned into fd-CAT2 for display were shown to bind to phOx-BSA (but not BSA) by ELISA (table 5). Phage were considered to be binding if A₄₀₅ of the sample was at least 10-fold greater than

the background in ELISA.

The phagemid vector, pHEN1 (fig. 26), is based upon pUC119 and contains restriction sites (SfiI and NotI) for cloning the fusion proteins. Here the transcription of antibody-g3p fusions is driven from the inducible lacZ promoter and the fusion protein targetted to the periplasm by means of the pelB leader. Phagemid was rescued with VCSM13 helper phage in 2xTY medium containing no glucose or IPTG: under these conditions there is sufficient expression of antibody-g3p. Fab and scFv fragments of NQ10.12.5 cloned into pHEN1 for display were shown to bind to pHx-BSA (but not BSA) by ELISA (Table 5) using the same criterion as above.

An alternative methodology for preparing libraries of Fab fragments expressed on the surface of phage would be to:

1. Prepare a library of phage expressing heavy chain (VHCH) genes from inserts in the phage genome.
 2. Prepare a library of light chain genes in a plamid expression vector in E.coli, preferably a phagemid, and isolate the soluble protein light chains expressed from this library.
 3. Bind the soluble protein light chains from the library to the heavy chain library displayed on phage.
 4. Select phage with the desired properties of affinity and specificity.
- These will encode the heavy chain (VHCH) genes.
5. Isolate the light chain genes encoding light chains which form suitable antigen binding sites in combination with the selected heavy chains, preferably by using superinfection of bacteria, containing phagemid expressing the light chain, with phage expressing the selected heavy chain (as described in example 20) and then assaying for antigen binding.

Example 26

Rescue of Phagemid Encoding a Gene III Protein Fusion with Antibody Heavy or Light Chains by Phage Encoding the Complementary Antibody Chain Displayed on Phage and the Use of this Technique to Make Dual Combinatorial Libraries

With random combinatorial libraries there is a limitation on the potential diversity of displayed Fab fragments due to the transformation efficiency of bacterial cells. Described here is a strategy (dual combinatorial libraries) to overcome this problem, potentially increasing the number of phage surveyed by a factor of 10^7 .

For assembly of heavy and light chains expressed from different vectors, phagemid (pHEN1-III or IV) was grown in E.coli HB2151 (a non-suppressor strain) to allow production of soluble chains, and rescued as above (example 27) except that helper phage were used

expressing partner chains as fusions to g3p (10^9 TU fd-CAT2-IV or III respectively) and 2 μ l tetracycline (12.5 mg/ml) in place of kanamycin.

5 Separate Vectors to Encode Fab Heavy and Light Chains

The heavy and light chains of Fab fragments can be encoded together in the same vector (example 25) or in different vectors. To demonstrate this the heavy chain (construct III) was cloned into pHEN1 (to provide soluble fragments) and the light chain (construct IV) into fd-CAT2 (to make the fusion with g3p). The phagemid pHEN1-III, grown in E.coli HB2151 (non-suppressor) was rescued with fd-CAT2-IV phage, and phage(mid) shown to bind to phOx:BSA, but not to BSA (Table 5). This demonstrates that soluble light chain is correctly associating with the heavy chain anchored to the g3p, since neither heavy chain nor light chain alone bind antigen (Table 5).

Similar results were obtained in the reverse experiment (with phagemid pHEN-1-IV and fd-CAT2-III phage) in which the heavy chain was produced as a soluble molecule and the light chain anchored to g3p (Table 5). Hence a Fab fragment is assembled on the surface of phage by fusion of either heavy or light chain to g3p, provided the other chain is secreted using the same or another vector (figure 28).

The resulting phage population is a mixture of phage abd rescued phagemid. The ratio of the two types of particle was assessed by infecting log phase E.coli TGI and plating on TYE plates with either 15 μ g/ml tetracycline (to select for fd-CAT2) or 100 μ g/ml ampicillin (to select for pHEN1). The titre of fd-CAT2 phage was 5×10^{11} TU/ml and the titre of pHEN1 2×10^{10} TU/ml, indicating a packaging ratio of 25 phage per phagemid.

Demonstrated here is an alternative strategy involving display of the heterodimeric antibody Fab fragments on the surface of phage. One of the chains is fused to g3p and the other is secreted in soluble form into the periplasmic space of the E.coli where it associates non-covalently with the g3p fusion, and binds specifically to antigen. Either the light or heavy chain can be fused to the g3p: they are displayed on the phage as Fab fragments and bind antigen (Figure 28). Described are both phage and phagemid vectors for surface display. Phagemids are probably superior to phage vectors for creation of large phage display libraries. Particularly in view of their higher transfection efficiencies (Two to three orders of magnitude higher), allowing larger libraries to be constructed. The phagemid vector, pHEN1 also allows the expression of soluble Fab fragments in non-suppressor E.coli.

Also demonstrated here is that heavy and light

chains encoded on the same vector (construct II), or on different vectors (constructs III and IV) can be displayed as Fab fragments. This offers two distinct ways of making random combinatorial libraries for display. Libraries of heavy and light chain genes, amplified by PCR, could be randomly linked by a 'PCR assembly' process (example 14) based on 'splicing by overlap extension', cloned into phage(mid) display vectors and expressed from the same promoter as part of the same transcript (construct II) as above, or indeed from different promoters as separate transcripts. Here the phage(mid) vector encodes and displays both chains. For a combinatorial library of 10^7 heavy chains and 10^7 light chains, the potential diversity of displayed Fab fragments (10^{14}) is limited by the transfection efficiency of bacterial cells by the vector (about 10^9 clones per μg cut and ligated plasmid at best) (W.J. Dower et al Nucl. Acids. Res. 16 6127-6145, 1988). Libraries thus prepared are analogous to the random combinatorial library method described by Huse, W.D. et al Science 246 1275-1281 (1989), but have the important additional feature that display on the surface of phage gives a powerful method of selecting antibody specificities from the large number of clones generated.

Alternatively, libraries of heavy and light chains could be cloned into different vectors for expression in the same cell, with a phage vector encoding the g3p fusion and a phagemid encoding the soluble chain. The phage acts as a helper, and the infected bacteria produced both packaged phage and phagemid. Each phage or phagemid displays both chains but encodes only one chain and thus only the genetic information for half of the antigen-binding site. However, the genes for both antibody chains can be recovered separately by plating on the selective medium, suggesting a means by which mutually complementary pairs of antigen binding heavy and light chain combinations could be selected from random combinatorial libraries. For example, a light chain repertoire on fd phage could be used to infect cells harbouring a library of soluble heavy chains on the phagemid. The affinity purified phagemid library could then be used to infect E.coli, rescued with the affinity purified phage library, and the new combinatorial library subjected to a further round of selection. Thus, antibody heavy and light chain genes are reshuffled after each round of purification. Finally, after several rounds, infected bacteria could be plated and screened individually for antigen-binding phage. Such 'dual' combinatorial libraries are potentially more diverse than those encoded on a single vector. By combining separate libraries of 10^7 light chain phage(mid)s, the diversity of displayed Fab fragments (potentially 10^{14}) is limited

only by the number of bacteria (10^{12} per litre). More simply, the use of two vectors should also facilitate the construction of 'hierarchical' libraries, in which a fixed heavy or light chain is paired with a library or partners (example 22), offering a means of 'fine-tuning' antibody affinity and specificity.

Example 27

Induction of Soluble scFv and Fab Fragments using Phagemid pHEN1

Further study of antibodies which have been expressed on the surface of phage would be greatly facilitated if it is simple to switch to expression in solution.

E. coli HB2151 was infected with pHEN phagemid (pHEN1-I or II), and plated on YTE, 100 µg/ml ampicillin plates. Colonies were shaken at 37°C in 2xTY medium, 100 µg/ml ampicillin, 1% glucose to $OD_{550}=0.5$ to 1.0. Cells were pelleted, washed once in 2xTY medium, resuspended in medium with 100 µg/ml ampicillin, 1 mM isopropyl β-D-thiogalactoside (IPTG), and grown for a further 16 hours. Cells were pelleted and the supernatant, containing the secreted chains, used directly in ELISA.

The phagemid pHEN1 has the advantage over phage fd-CAT2, in that antibody can be produced either for phage display (by growth in supE strains of *E. coli*) or as a tagged soluble fragment (by growth in non-suppressor strains), as a peptide tag (example 24) and amber codon were introduced between the antibody and g3p. Secretion of soluble Fab fragments from pHEN1-II or scFv fragments from pHEN1-I was demonstrated after growth in *E. coli* HB2151 and induction with IPTG using Western blots (Figure 29). For detection of secreted proteins, 10 µl supernatant of induced cultures were subjected to SDS-PAGE and proteins transferred by electroblotting to Immobilon-P (Millipore). Soluble heavy and light chain were detected with goat polyclonal anti-human Fab antiserum (Sigma) and peroxidase conjugated rabbit anti-goat immunoglobulin (Sigma), each at a dilution of 1:1000. The tagged VK domain was detected with 9E10 antibody (1:1000) and peroxidase conjugated goat anti-mouse immunoglobulin (Fc specific) (1:1000) (Sigma) or with a peroxidase labelled anti-human CK antiserum (Dako). 3,3'-diaminobenzidine (DAB; Sigma) was used as peroxidase substrate (Harlow E., et al. 1988 Supr). With the scFv, the fragments were detected using the 9E10 anti-myc tag antibody (data not shown). With the Fab, only the light chain was detected by 9E10 (or anti-human CK) antibody, as expected, while the anti-human Fab antiserum detected both heavy and light chains. Binding of the soluble scFv and Fab fragments to phOx-BSA (but not to BSA) was also demonstrated by ELISA (Table 5B). Thus scFv and Fab fragments can be displayed on phage or

secreted as soluble fragments from the same phagemid vector.

Example 28

Increased Sensitivity in ELISA assay of Lysozyme using FDTscFvD1.3 as Primary Antibody Compared to Soluble scFvD1.3

In principle the use of phage antibodies should allow more sensitive immunoassays to be performed than with soluble antibodies. Phage antibodies combine the ability to bind a specific antigen with the potential for amplification through the presence of multiple (ca.2800) copies of the major coat protein (g8p) on each virion. This would allow the attachment of several antibody molecules directed against M13 to each virion followed by the attachment of several molecules of peroxidase-conjugated anti-species antibody (anti-sheep) IgG in the case below). Thus for every phage antibody bound to antigen there is the potential for attaching several peroxidase molecules whereas when a soluble antibody is used as the primary antibody this amplification will not occur.

ELISA plates were coated overnight at room temperature using 200 μ l of 10 fold dilutions of hen egg lysozyme (1000, 100, 10, 1, 0.1 and 0.01 μ g/ml) in 50mM NaHCO₃, pH9.6. ELISA was performed as described in example 4 except that (i) incubation with anti-lysozyme antibody was with either FDTscFvD1.3 (pAb; 10^{11} phage per well; 1.6mol) or soluble affinity purified scFvD1.3 (18 μ g per well; 0.7nmol) (ii) incubation with second antibody was with 1/100 dilution of sheep anti-M13 serum for FDTscFvD1.3 samples or with or 1/100 dilution of rabbit anti-scFvD1.3 serum (from S. Ward) for soluble scFvD1.3 samples (iii) peroxidase-conjugated rabbit anti-goat immunoglobulin (Sigma; 1/5000) was used for FDTscFvD1.3 samples and peroxidase-conjugated goat anti-rabbit immunoglobulin (Sigma; 1/5000) was used for soluble scFvD1.3 samples. Absorbance at 405nm was measured after 15h. The results are shown in Figures 30 and 31. In these figures lysozyme concentrations for coating are shown on a log scale of dilutions relative to 1 μ g/ml. (i.e. log = -3 = 1mg/ml ; log = 2 = 0.01 μ g/ml)

Higher signals were obtained with FDTscFvD1.3 at all concentrations of lysozyme (Fig.31) but the difference was very marked at the greatest dilutions, where antigen quantities are most limiting (Figs. 30 and 31). This suggests that phage antibodies may be particularly valuable for sandwich type assays where the capture of small amounts of antigen by the primary antibody will generate an amplified signal when phage antibodies directed against a different epitope are used as the second antigen binding antibody.

Example 29

Direct Rescue and Expression of Mouse Monoclonal Antibodies as Single Chain Fv Fragments on the Surface of Bacteriophage fd.

5 The principle is very similar to that described in example 14. It consists of the PCR assembly of single chain antibodies from cDNA prepared from mouse monoclonals. As an example, the rescue and expression of two such antibodies from monoclonals expressing antibodies against the steroid hormone oestriol is described.

10 A. RNA Preparation

RNA can be prepared using many procedures well known to those skilled in the art. In this example, the use of Triton X-100 lysis, phenol/SDS RNase inactivation gave excellent results.

1. The mouse monoclonal cells that were used here had been harvested by centrifugation and resuspended in serum free medium. They were then centrifuged and resuspended in saline and after a final centrifugation step, resuspended in sterile water at 1×10^7 cells per ml. (Normally cells would be washed in PBS buffer and finally resuspended in PBS buffer, but these particular cells were supplied to us as described frozen in water.).

2. To 750ul of cells was added 250ul of ice cold 4X lysis buffer (40mM Tris HCl pH 7.4/4mM $MgCl_2$ /600mM NaCl/40mM VRC (Veronyl ribosyl complex)/2% Triton X-100). The suspension was mixed well and left on ice for 5 minutes.

3. Centrifugation was carried out at 4°C in a microfuge at 13000 rpm for 5 min.

The supernatant is then phenol extracted three times, phenol chloroform extracted three times and finally, ethanol precipitated as described in the materials and methods. The precipitate was resuspended in 50ul water.

35 4. The optical density of the RNA at 260nm with a 2.5ul sample in 1ml water was measured. The RNA was checked by electrophoresis of a 2ug sample on a 1% agarose gel. RNA in the range of 32ug to 42ug was obtained by this method.

40 B. cDNA Preparation

The method used is the same as that described in example 14. Two cDNA preparations were made. These were from RNA extracted from the monoclonals known as cell lines 013 and 014 which both express antibodies against eh steroid hormone, oestriol.

45 C. Primary PCRs

50 The method used is essentially the same as that described in example 14. The VH region was amplified with the primers VH1BACK and VH1FOR-2. For the Vkappa region, four separate reactions were carried out using the primer VK2BACK and wither MJK1FONX, MJK2FONX, MJK4FONX or MJK5FONX. Samples (5ul) were checked on a 1.5% agarose gel. From this it was observed that for

cDNA prepared from the two oestriol monoclonals the primers VK2BACK and MJK1FONX gave the best amplification of the Vkappa region. The VH bands and the Vkappa bands amplified with VK2BACK/MJK1FONX were purified on 2% low melting point agarose gels for each monoclonals. The DNA bands were excised from the gel and purified using a dedicated Geneclean kit as described in example 14.

D. Preparation of linker

The method used is essentially the same as that described in example 14. In this case, the amplified linker DNA was purified on a 2% agarose gel and recovered from the gel with a dedicated "Mermaid" kit (BIO 101, Geneclean, La Jolla, San Diego, California, USA) using the manufacturers instructions.

E. Assembly PCRs

The method used is essentially the same as that described in example 14. In this case, the assembled PCR product was purified on a 2% agarose gel and recovered from the gel with a dedicated "Mermaid" kit.

F. Adding restriction sites and work-up

The assembled product was "tagged" with Apa LI and Not I restriction sites. The DNA was then digested with Apa LI and Not I to give the appropriate sticky ends for cloning and then purified on a 2% low melting point agarose gel and extracted using a Geneclean kit. The method used is the same as that described in example 14.

G. Cloning into Vector fd-CAT2

A total of 15ug of CsCl purified fd-CAT2 DNA was digested with 100 units of the restriction enzyme Not I (New England Biolabs) in a total volume of 200ul 1X NEB Not I buffer with 1X NEB acetylated BSA for a total of 3 hours at 37°C. The vector DNA was then treated twice with 15ul Strataclean (a commercially available resin for the removal of protein), following the manufacturers instructions (Stratagene, 11099 North Torrey Pines Road, La Jolla, California, USA). The DNA was then ethanol precipitated and redissolved in TE buffer (Sambrook et al., 1989 supra). The DNA was then digested with 100 units of the restriction enzyme Apa LI (New England Biolabs) in a total volume of 200ul 1X NEB Buffer 4 overnight at 37°C. The vector was then purified with a Chroma Spin 1000 column following the manufacturers instructions (Clontech Laboratories Inc, 4030 Fabian way, Palo Alto, California, USA). This step removes the Apa LI/Not I fragment to give cut vector DNA for maximum ligation efficiency.

Ligation reactions were carried out with 2.5-10ng of the DNA insert and 10ng of vector in a total volume of 10ul of 1X NEB ligase buffer with 1ul of NEB ligase (New England Biolabs) at 16°C overnight (approx 16 hours).

H. Transformation and growth

E.coli strain TGI was made competent and transformed

with the fdCAT2 recombinant DNA as described by Sambrook et al, 1989 Supra. The cells were plated out on LBtet plates (10g tryptone, 5g yeast extract, 10g NaCl, 15g bacto-agar per litre with 15ug/ul of tetracycline added just before pouring the plates) and grown overnight.

Single well isolated colonies were then inoculated into 10 ml of LBtet broth (LB medium with 15ug/ul of tetracycline) in 50 ml tubes. After overnight growth at 35°C/350rpm in a bench top centrifuge. The supernatants were transferred to 15 ml centrifuge tubes and 2ml 20% PEG 8000/2.5M NaCl added to each. After incubating at room temperature for 20-30 minutes, the recombinant phage was pelleted by centrifugation at 9000rpm in a Sorval SM24 rotor for 30 minutes. The PEG supernatant was discarded. Any remaining PEG was removed with a pasteur pipette after a brief (2 minutes) centrifugation step. This last step was repeated to make sure that no PEG remained. The phage pellet was then resuspended in 500ul PBS buffer. This was transferred to a microcentrifuge tube and spun at 13000 rpm to remove any remaining cells. The phage supernatant was transferred to a fresh tube.

I. Assay for antibody expression

Bacteriophage fd recombinants were screened for the expression of antibody against oestriol by ELISA. This method is described in example 6. In this case the following alterations are relevant.

1. Microtitre plates were coated overnight with 40ug/ml oestriol-6 carboxymethylloxime-BSA (Steraloids, 31 Radcliffe Road, Croydon, CR0 5QJ, England).
2. 1st antibody was the putative phage anti oestriol antibody. 50ul of phage in a final volume of 200ul of sterile PBS combining 0.25% gelatin was added to each well.
3. 2nd antibody was sheep anti M13 at 1:1000 dilution.
4. 3rd antibody was peroxidase conjugated rabbit anti goat immunoglobulin.

Recombinants expressing functional antibody were detected by incubation with the chromogenic substrate 2'2' axinobis (3-ethyl benzthiazoline sulphonic acid). The results are shown in figures 32 and 33.

Example 30

Kinetic Properties of Alkaline Phosphatase Displayed on the Surface of Bacteriophage fd

This example demonstrates that kinetic properties of an enzyme expressed on phage are qualitatively similar to those in solution. Bacteriophage fd displaying alkaline phosphatase fusions of gene 3 with either the native arginine (see example 31) or the mutant residue alanine at position 166 (see example 11) were prepared by PEG precipitation as described in the materials and methods.

The kinetic parameters of alkaline phosphatase expressed on the surface of fd phage were investigated in

1M Tris/HCl, pH8.0 at 20°C with 1ml 4-nitrophenyl phosphate as substrate. The reactions were initiated by the addition of 100µl of a phage-alkaline phosphatase fusion preparation, 50 fold concentrated with respect to the original culture supernatant. The rate of change of absorbance was monitored at 410nm using a Philips 8730 spectrophotometer and the initial reaction rate calculated using a molar absorbance of 16200 l/mol/cm. For the fdphoAla 166 enzyme but not fdphoArg166 a lag phase was seen following this addition, the reaction rate accelerating until a steady state was obtained after approximately 60 to 90 secs. This steady state rate was used for determination of kinetic parameters. No deviation from Michaelis Menten kinetics was apparent for either phage enzyme. Values of K_m and k_{cat} were derived from plots of s/v against s and are shown in Table 6.

Because of the difficulty in establishing the relationship between the number of phage particles and the number of active enzyme dimers formed on the phage k_{cat} values are expressed not as absolute values, but as relative values between the two enzyme forms. Western blots (carried out as in example 31 using antiG3p antiserum) of the phage enzyme preparations used in this experiment showed approximately equal intensities for the full length fusion band with the Arg166 and Ala166 enzymes when detected using antibody directed against gene3. In these preparations the intact fusion represents approximately 30% of the detected material. The two preparations were therefore assumed to be expressing approximately the same concentrations of intact fusions.

Table 6 summarises the kinetic data from this experiment and compares it with data from Chaidaroglou, A. et al (Biochemistry 27, 8338-8343 (1988)) obtained with soluble preparations of the wild type and mutant enzyme forms. The same substrate and assay conditions were used in both experiments. Soluble alkaline phosphatase was also tested in parallel in our experiments ($K_m=8.5\mu M$; $k_{cat}=3480$ mol substrate converted mol enzyme⁻¹ min⁻¹).

The effect of mutating arginine at position 166 to alanine is qualitatively similar for the phage enzyme as for the soluble enzyme. K_m is increased about 15 fold and the relative k_{cat} is decreased to 36% of that for wild type. This increased K_m would reflect a reduction in substrate affinity in the phage enzyme on mutation of Arg166, as was proposed for the soluble enzyme (Chaidaroglou et al, 1988 supra), assuming the same kinetic mechanism applies. There are, however, some quantitative differences in the behaviour of K_m of the phage enzyme. The K_m of 73µM observed for fdphoArg166 compares with a K_m of 12.7µM for the free enzyme; the K_m

for fdphoAla166 is 1070 μ M whereas the free mutant enzyme has a K_m of 1620 μ M. One can speculate that the higher K_m for fdphoArg 166 and the lower K_m for fdphoAla166, compared to the soluble enzymes result from the 'anchored' alkaline phosphatase fusion molecules interacting to form dimers in a different manner to the enzyme in free solution.

The relative values of k_{cat} for the Arg166 and Ala166 forms are however very similar for both the phage enzymes and the soluble enzymes, a reduction occurring on mutation to 35 to 40% of the value for the native enzyme. The rate limiting step, determining k_{cat} , for soluble phoArg166 is thought to be dissociation of non-covalently bound phosphate from the enzyme (Hull W.E. et al. Biochemistry 15, 1547-1561 1976). Chaidaroglou et al (1988) supra suggest that, for the soluble enzyme, mutation of Arg166 to alanine alters additional steps, one of which may be hydrolysis of the phosphoenzyme intermediate. The similarity in the reduction in k_{cat} on mutation of Arg166 to alanine for the phage enzymes suggests that the same steps may be altered in a quantitatively similar manner in the mutant phage enzyme as in the mutant soluble enzyme.

Thus, enzymes displayed on phage show qualitatively similar characteristics to soluble enzymes.

Example 31

Demonstration using Ultrafiltration that Cloned Alkaline Phosphatase Behaves as Part of the Virus Particle

The construct fdphoAla166 (derived in example 11) was converted back to the wild type residue (arginine) at position 166 by in vitro mutagenesis (Amersham International) using the primer APARG166:5' TAGCATTTGCGCGAGGTCACA 3'.

This construct with the wild type insert was called fdphoArg166.

E.coli TG1 or KS272 cells (cells with a deletion in the endogenous phoA gene, Strauch and Beckwith, 1988 Supra) containing either fd-phoAla166, fdphoArg166 or fd-CAT2 were grown for 16 hours at 37°C in 2xTY with 15 μ g/ml tetracycline. Concentrated phage were prepared as follows. Phage-enzyme cultures are clarified by centrifugation (15 min at 10,000 rpm, 8 x 50 ml rotor, sorval RC-5B centrifuge). Phage are precipitated by adding 1/5 volume 20% polyethylene glycol, 2.5 M NaCl, leaving for 1 hr at 4°C, and centrifuging (as above). Phage pellets are resuspended in 10 mM Tris-HCl, pH 8.0 to 1/100th of the original volume, and residual bacteria and aggregated phage removed by centrifugation for 10 to 15 minutes in a bench microcentrifuge at 13000 rpm at 4°C.

SDS/Polyacrylamide gel electrophoresis and western blotting were basically as described previously (example

2). Denatured samples consisting of 16µl of a 50 fold concentrate of phage were separated using a 10% SDS/polyacrylamide gel and detected with polyclonal antiserum raised against either E.coli alkaline phosphatase (Northumbria Biologicals, South Nelson Industrial Estate, Cramlington, Northumberland, NE23 9HL) or against the minor coat protein encoded by gene 3 (from Prof. I. Rasched, Universitat Konstanz, see Stengele et al, 1990) at 1 in 1000 dilution. This was followed by incubation with peroxidase-conjugated goat-anti-rabbit immunoglobulin (Sigma 1 in 5000) and detection with the ECL Western blotting system (Amersham International).

The presence of fusion proteins was confirmed by western blotting of proteins from phage particles derived from fd-phoAla166 (phage-enzyme) or fd-CAT2 (vector phage). Detection with antiserum raised against the gene 3 protein reveals a product of apparent relative molecular mass (Mr) of 63,000 in vector phage (figure 34e). Although this is different from the predicted molecular weight based on the amino acid sequence (42,000), the natural product of gene 3 has previously been reported to exhibit reduced mobility during electrophoresis (Stengele et al, 1990).

In the fd-phoAla166 sample the largest band has an apparent Mr of 115,000, (fig. 34). Taking into account the aberrant mobility of the gene 3 portion of the fusion, this is approximately the size expected from fusing with an alkaline phosphatase domain of 47 kD. This analysis also reveals that a proportion of the Gene3 reactive material in this phage-enzyme preparation is present at the size of the native gene3 product, suggesting that degradation is occurring. In the preparation shown in figure 34, approximately 5-10% of the gene 3 fusions are intact. In more recent preparations and in all the preparations used in this example and example 32, approximately 30-60% of fusions are full length.

The protein of Mr 115,000 is the major protein observed in Western blots of phage-enzyme derived from TG1 cells when probed with antiserum raised against E.coli alkaline phosphatase (anti-BAP), confirming the assignment of this band to intact fusion. Further, when phage enzyme is prepared using KS272 cells, which have a deletion in the endogenous phoA gene (Strauch & Beckwith, 1988, supra.) it is also the major band. There are additional bands at Mr 95000 and 60000 reactive with anti-BAP antiserum which may indicate degradation of the fusion product.

The anti-BAP antiserum also reacts with material running with the dye front and with a molecule of Mr 45,000 but evidence suggests that this material is not alkaline phosphatase. This pattern is detected in PEG

precipitated vector phage samples (figure 34c) and is not therefore contributed by protein expressed from the cloned *phoA* gene. These bands are detected in culture supernatants of cells carrying fd-CAT2 but is not detected in the supernatant of uninfected cells (not shown) and so either represents cross-reactivity with phage encoded material or with a PEG precipitable cellular component leaked from infected cells (Boeke et al, Mol. Gen. Genet. 186, 185-192 1982). Although the fragment of Mr, 45,000 is close to the size of free alkaline phosphatase (47,000), it is present in phage preparations from KS272 cells which have a deletion in the *phoA* locus. Furthermore its mobility is different from purified alkaline phosphatase and they can be distinguished by electrophoresis (figure 34d).

Ultrafiltration was used to confirm that the fusion protein behaved as though it were part of a larger structure, as would be expected for an enzyme bound to a phage particle. Phage samples (100 μ l of a 50 fold concentrate) were passed through ultrafiltration filters with a nominal molecular weight limit of 300000 daltons (Ultrafree-MC filters, Millipore) by centrifugation for 5 to 15 minutes at 13,000 r.p.m. in an MSE microcentaur microfuge. Retained material was recovered by resuspending in 100 μ l of 10mM Tris, pH 8.0.

Phage-enzyme or free alkaline phosphatase (83ng) mixed with vector phage were passed through filters with a nominal molecular weight limit of 300,000 daltons (Ultrafree-MC filters, Millipore). Figure 35 A again shows that the band of Mr, 115,000 is the major product reactive with anti-BAP antiserum. This and the other minor products reactive with anti-BAP are present in material retained by the ultrafiltration membrane. Analysis of retained and flow through fractions of phage preparations derived from KS272 demonstrates that different molecular species are being separated by the ultrafiltration membranes. Figure 35b shows the protein of Mr 115,000 is retained by the filter whereas the putative degradation products of Mr 95,000 and 60,000 found in phage preparations derived from KS272 cells, are not retained.

In mixture of alkaline phosphatase and vector phage Figure 35c-f, free alkaline phosphatase (dimer size of 94,000 daltons) is detected in the flow through as a monomer band with Mr 47,000 on denaturing polyacrylamide gels (figure 35B), while the cross reactive molecule found in vector phage preparations (Mr 45,000) is retained on the filter (figure 35B). This suggests that the cross reactive molecule is part of the phage particle and underlines the fact that the ultrafiltration membranes are effecting a separation. Thus the expected fusion band in this phage-enzyme is present in material

retained on ultrafiltration membranes demonstrating that it is part of a larger structure as would be expected for viral bound enzyme.

5 Catalytic activity has been demonstrated on phage particles expressing alkaline phosphatase. Table 7 shows that the wild type alkaline phosphatase gene expressed on phage (fd-phoArg166) has a specific activity (moles of substrate converted per mole of viral particles) of 3,700/min. This is close to the turnover value of 10 4540/min found for purified alkaline phosphatase by Malamy and Horecker, Biochemistry 3, 1893-1897 1964).

Chaidaroglou et al, 1988 supra have shown that substituting alanine for arginine at the active site (residue 166) leads to a reduction in the rate of 15 catalysis. Preparations of phage displaying alkaline phosphatase with this mutation derived from TG1 and KS272 show reduced specific activities of 380 and 1400 mol substrate converted/mol phage/min respectively. Enzyme activity was measured in the retained and flow-through 20 fractions prepared by ultrafiltration, shown in figure 35. The bulk of activity from phage-enzyme was retained on the filters whereas the majority of activity from free enzyme passes through. Therefore, the enzyme activity in these fusions behaved as would be expected for virally 25 associated enzyme (not shown). Little or no catalytic activity is measured in preparations of vector phage from either TG1 or KS272 cells (Table 7), indication that the catalytic activities above are due to phage enzyme and not contamination with bacterial phosphatase. Addition 30 of phage particles to soluble enzyme does not have a significant effect on activity (Table 7).

Therefore, both the catalytic and immunochemical activity of alkaline phosphatase have been demonstrated to be due to enzyme which is part of the phage particle.

35 Example 32

Affinity chromatography of phage alkaline phosphatase

Affinity chromatography, using the specific binding properties of enzymes has proved to be a very powerful method for their purification. The purification of 40 phage-enzymes by this approach would enable the genetic material encoding the enzyme to be isolated with the enzyme itself. Thus, mutagenesis of cloned enzymes expressed on the surface of filamentous bacteriophage will lead to a whole population of enzyme variants, from 45 which variants with desired binding properties could be isolated.

Soluble alkaline phosphatase (from calf intestine) has been purified by binding to immobilised arsenate (a competitive inhibitor), and eluting with inorganic 50 phosphate, which is a product (and competitive inhibitor) of the enzyme reaction (Brenna, O. et al, Biochem. J. 151 291-296 1975). The applicants have determined that